

FORM PTO-1390 (Modified) (REV. 11-2000)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER 235.00270101	
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371				U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR 10/018936	
INTERNATIONAL APPLICATION NO. PCT/US00/10672		INTERNATIONAL FILING DATE 20 April 2000		PRIORITY DATE CLAIMED 21 April 1999	
TITLE OF INVENTION CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS					
APPLICANT(S) FOR DO/EO/US Salman BAIG, Raymond T. DAMIAN, Arthur Clinton WHITE					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:					
1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input checked="" type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (24) indicated below. 4. <input checked="" type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (Article 31). 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371 (c) (2)) a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> has been communicated by the International Bureau. c. <input checked="" type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). a. <input type="checkbox"/> is attached hereto. b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4). 7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3)) a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> have been communicated by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input checked="" type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)). 10. <input type="checkbox"/> An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)). 11. <input checked="" type="checkbox"/> A copy of the International Preliminary Examination Report (PCT/IPEA/409). 12. <input checked="" type="checkbox"/> A copy of the International Search Report (PCT/ISA/210).					
Items 13 to 20 below concern document(s) or information included:					
13. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 14. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 15. <input checked="" type="checkbox"/> A FIRST preliminary amendment. 16. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 17. <input type="checkbox"/> A substitute specification. 18. <input type="checkbox"/> A change of power of attorney and/or address letter. 19. <input checked="" type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825. 20. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4). 21. <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4). 22. <input checked="" type="checkbox"/> Certificate of Mailing by Express Mail 23. <input checked="" type="checkbox"/> Other items or information:					
Written Sequence Listing, Return itemized postcard, transmittal letter, and Communication Regarding Entry of Sequence Listing					

U.S. APPLICATION NO. (PCT/US/00/10672)		ATTORNEY'S DOCKET NUMBER 235.00270101	
24. The following fees are submitted:		CALCULATIONS PTO USE ONLY	
BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) : <input type="checkbox"/> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1040.00 <input checked="" type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$890.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$740.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$710.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00			
ENTER APPROPRIATE BASIC FEE AMOUNT =		\$890.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than months from the earliest claimed priority date (37 CFR 1.492 (e)). <input type="checkbox"/> 20 <input type="checkbox"/> 30		\$0.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	48 - 20 =	28	x \$18.00
Independent claims	12 - 3 =	9	x \$84.00
Multiple Dependent Claims (check if applicable).			<input type="checkbox"/> \$0.00
TOTAL OF ABOVE CALCULATIONS =			\$2,150.00
<input checked="" type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.			\$1,075.00
SUBTOTAL =			\$1,075.00
Processing fee of \$130.00 for furnishing the English translation later than months from the earliest claimed priority date (37 CFR 1.492 (f)). <input type="checkbox"/> 20 <input type="checkbox"/> 30 +			\$0.00
TOTAL NATIONAL FEE =			\$1,075.00
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable). <input type="checkbox"/>			\$0.00
TOTAL FEES ENCLOSED =			\$1,075.00
			Amount to be: refunded \$
			charged \$
a. <input type="checkbox"/> A check in the amount of _____ to cover the above fees is enclosed. b. <input checked="" type="checkbox"/> Please charge my Deposit Account No. <u>13-4895</u> in the amount of <u>\$1,075.00</u> to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>13-4895</u> A duplicate copy of this sheet is enclosed. d. <input type="checkbox"/> Fees are to be charged to a credit card. WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.			
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.			
SEND ALL CORRESPONDENCE TO: <div style="border: 1px solid black; padding: 5px; width: 40%;"> Victoria A. Sandberg Mueeting, Raasch & Gebhardt, P.A. P.O. Box 581415 Minneapolis, Minnesota 55458-1415 United States of America </div> <div style="width: 60%;"> SIGNATURE <u>Victoria A. Sandberg</u> NAME <u>Victoria A. Sandberg</u> 41,287 REGISTRATION NUMBER October 19, 2001 DATE </div>			

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant(s): Salman Baig et al.) Group Art Unit: Unknown
Serial No.: Unknown) Examiner: Unknown
(U.S. National Stage of)
PCT/US00/10672))
Filed: Herewith)
(Int'l. Filing Date)
April 20, 2000))

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND
TREATMENT OF NEUROCYSTICERCOSIS

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
ATTN: BOX PCT
Washington, D.C. 20231

Dear Sir:

The present application is a U.S. National Stage of PCT Application Serial No. PCT/US00/10672 filed April 20, 2000, which claims priority to U.S. Provisional Patent Application Serial No. 60/130,338 filed April 21, 1999.

Prior to taking up the above-identified application for examination, please amend the application as follows:

In the Title

Please replace the title with:

"CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND
TREATMENT OF NEUROCYSTICERCOSIS"

In the Claims

Please amend claims 4-6, 8-10, 14, 15, 17-25, 28, 29, 31-34, 39, 42 and 43, and add claims 44-48. The new and amended claims are provided below in clean form. Per 37 C.F.R. §1.121, amended claims are also shown in Appendix A with notations to indicate changes made

Preliminary Amendment

Page 2

Applicant(s): Salman Baig et al.

Serial No. Unknown (National Stage of PCT/US00/10672)

Filed: Herewith (Int'l. Filing Date: April 20, 2000)

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF
NEUROCYSTICERCOSIS

(for convenience, all pending claims, including those added hereby, are provided in Appendix A).

4. (AMENDED) A vaccine comprising a cyst wall cysteine proteinase or a polynucleotide having a nucleic acid sequence encoding a cyst wall cysteine proteinase; and a pharmaceutically acceptable carrier.

5. (AMENDED) A method for preparing the vaccine of claim 4 comprising mixing a cyst wall cysteine proteinase or a polynucleotide having a nucleic acid sequence encoding a cyst wall cysteine proteinase; and a pharmaceutically acceptable carrier.

6. (AMENDED) A vaccine comprising at least one component selected from the group consisting of (a) a polypeptide comprising a cyst wall cysteine proteinase or an immunogenic polypeptide subunit thereof and (b) a polynucleotide comprising a nucleotide sequence encoding a polypeptide comprising a cyst wall cysteine proteinase or an immunogenic polypeptide subunit thereof; and a pharmaceutically acceptable carrier.

8. (AMENDED) A method for treating an animal harboring a *Taenia* infection comprising administering to the infected animal the vaccine of claim 6, wherein administration of the vaccine is effective to eliminate the parasite from the animal or to prevent or delay the appearance of cysticercosis or neurocysticercosis the animal.

9. (AMENDED) A method for protecting an animal against a *Taenia* infection comprising administering to an uninfected animal the vaccine of claim 6, wherein administration of the composition is effective to prevent subsequent infection of the animal by the parasite or to prevent the development of cysticercosis or neurocysticercosis in the animal after subsequent infection by the parasite.

Preliminary Amendment

Page 3

Applicant(s): Salman Baig et al.

Serial No. Unknown (National Stage of PCT/US00/10672)

Filed: Herewith (Int'l. Filing Date: April 20, 2000)

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF
NEUROCYSTICERCOSIS

10. (AMENDED) The method of claim 8 wherein the animal is a pig or a human and the *Taenia* infection is a *T. solium* infection.

14. (AMENDED) The pharmaceutical composition of claim 12 wherein R comprises a chemical moiety selected from the group consisting of a carboxylic acid derivative, an amide derivative, a benzene derivative, a phenyl derivative, a chloromethylketone or derivative thereof, a fluoromethylketone or derivative thereof, an alphaketo acid or derivative thereof, a ketoamide or derivative thereof, a ketoester or derivative thereof, a vinylsulfone or derivative thereof, and a pyridyl or derivative thereof.

15. (AMENDED) The pharmaceutical composition of claim 12 wherein R comprises a fluoromethylketone (FMK).

17. (AMENDED) The pharmaceutical composition of claim 11 wherein the inhibitor molecule inhibits cyst wall cysteine proteinase derived from *T. solium* or *T. crassiceps*.

18. (AMENDED) A pharmaceutical composition comprising Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK; and a pharmaceutically acceptable carrier.

19. (AMENDED) A method for treating human neurocysticercosis comprising administering to a human subject the pharmaceutical composition of claim 18.

20. (AMENDED) A method for treating porcine cysticercosis comprising administering to a porcine subject the pharmaceutical composition of claim 18.

Preliminary Amendment

Page 4

Applicant(s): Salman Baig et al.

Serial No. Unknown (National Stage of PCT/US00/10672)

Filed: Herewith (Int'l. Filing Date: April 20, 2000)

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF
NEUROCYSTICERCOSIS

21. (AMENDED) A method for treating human neurocysticercosis or porcine cysticercosis comprising administering to a human or porcine subject a pharmaceutical composition comprising an inhibitor molecule that inhibits the activity of a *Taenia* cysteine protease, the inhibitor molecule comprising a peptide or peptidomimetic compound; and a pharmaceutically acceptable carrier.
22. (AMENDED) The method of claim 21 wherein the peptide or peptidomimetic compound comprises (Xaa)_n-Yaa-Zaa-R; wherein Xaa and Zaa are each independently any amino acid; Yaa is a hydrophobic amino acid; R comprises a nucleophilic moiety; and n = 0-5.
23. (AMENDED) The method of claim 22 wherein Yaa is leucine.
24. (AMENDED) The method of claim 22 wherein R comprises a chemical moiety selected from the group consisting of a carboxylic acid derivative, an amide derivative, a benzene derivative, a phenyl derivative, a chloromethylketone or derivative thereof, a fluoromethylketone or derivative thereof, an alphaketo acid or derivative thereof, a ketoamide or derivative thereof, a ketoester or derivative thereof, a vinylsulfone or derivative thereof, and a pyridyl or derivative thereof.
25. (AMENDED) The method of claim 22 wherein R comprises a fluoromethylketone (FMK).
28. (AMENDED) The method of claim 26 wherein R comprises a chemical moiety selected from the group consisting of a carboxylic acid derivative, an amide derivative, a benzene derivative, a phenyl derivative, a chloromethylketone or derivative thereof, a fluoromethylketone or derivative thereof, an alphaketo acid or derivative thereof, a ketoamide or derivative thereof, a ketoester or derivative thereof, a vinylsulfone or derivative thereof, and a pyridyl or derivative thereof.
29. (AMENDED) The method of claim 26 wherein R comprises a fluoromethylketone (FMK).

Preliminary Amendment

Page 5

Applicant(s): Salman Baig et al.

Serial No. Unknown (National Stage of PCT/US00/10672)

Filed: Herewith (Int'l. Filing Date: April 20, 2000)

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF
NEUROCYSTICERCOSIS

31. (AMENDED) The method of claim 26 performed in cell free environment.
32. (AMENDED) The method of claim 26 performed in cell culture, in an organ, or in a tissue.
33. (AMENDED) The method of claim 26 performed in a whole animal.
34. (AMENDED) The method of claim 26 wherein the *Taenia* cysteine proteinase is derived from *T. solium*.
39. (AMENDED) The method of claim 37 wherein the inhibitor is Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK.
42. (AMENDED) The method of claim 40 wherein inhibitor is Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK.
43. (AMENDED) The method of claim 40 wherein the animal is a pig or a human and the *Taenia* infection is a *T. solium* infection.
44. (NEW) The method of claim 9 wherein the animal is a pig or a human and the *Taenia* infection is a *T. solium* infection.
45. (NEW) The method of claim 38 wherein the inhibitor is Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK.
46. (NEW) The method of claim 41 wherein inhibitor is Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK.

Preliminary Amendment

Page 6

Applicant(s): Salman Baig et al.

Serial No. Unknown (National Stage of PCT/US00/10672)

Filed: Herewith (Int'l. Filing Date: April 20, 2000)

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF
NEUROCYSTICERCOSIS

47. (NEW) The method of claim 41 wherein the animal is a pig or a human and the *Taenia* infection is a *T. solium* infection.

48. (NEW) The method of claim 42 wherein the animal is a pig or a human and the *Taenia* infection is a *T. solium* infection.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
217

Preliminary Amendment

Page 7

Applicant(s): Salman Baig et al.

Serial No. Unknown (National Stage of PCT/US00/10672)

Filed: Herewith (Int'l. Filing Date: April 20, 2000)

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

Remarks

Claims 4-6, 8-10, 14, 15, 17-25, 28, 29, 31-34, 39, 42 and 43 having been amended and claims 44-48 have been added, the pending claims are 1-48.

Amendments to the claims have been made to remove multiple dependency, and to clarify the claimed subject matter.

Conclusion

The Examiner is invited to contact Applicants' Representatives at the below-listed telephone number, if there are any questions regarding this Preliminary Amendment or if prosecution of this application may be assisted thereby.

Respectfully submitted for

Salman Baig et al.

By

Mueting, Raasch & Gebhardt, P.A.

P.O. Box 581415

Minneapolis, MN 55458-1415

Phone: (612)305-1220

Facsimile: (612)305-1228

Customer Number 26813

By: 

Victoria A. Sandberg

Reg. No. 41,287

Direct Dial (612) 305-1226

October 19, 2001

Date

"Express Mail" mailing label number EL88271991US

Date of Deposit October 19, 2001

I hereby certify that this paper and /or fee is/are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, AIA: Box PCT, Washington, D. C. 20231.

Name: Gara L. Ludwig
Gara L. Ludwig

9/PRTS

531 Rec'd PCT. 10/018936
19 DEC 2001

APPENDIX A - SPECIFICATION/CLAIM AMENDMENTS
INCLUDING NOTATIONS TO INDICATE CHANGES MADE
Serial No.: Unknown (National Stage of PCT/US00/10672)
Docket No. 235.0027 0101

Amendments to the following are indicated by underlining what has been added and bracketing what has been deleted. Additionally, all amendments have been shaded.

In The Title

The title has been amended as follows:

CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND
TREATMENT OF ~~NEUROCYSTICERCOSIS~~ ~~NEUROCYSTICEROCOSIS~~

In the Claims

For convenience, all pending claims are shown below.

1. An isolated cyst wall cysteine proteinase derived from *Taenia solium*.
2. An isolated polynucleotide having a nucleic acid sequence that encodes the proteinase of claim 1.
3. An isolated polynucleotide that is the complement of the nucleic acid molecule of claim 2.
4. (AMENDED) A ~~vaccine comprising a~~ cyst wall cysteine proteinase or a polynucleotide having a nucleic acid sequence encoding a cyst wall cysteine proteinase, ~~and a pharmaceutically acceptable carrier~~, for use as a vaccine.
5. (AMENDED) ~~A method for preparing the vaccine of claim 4 comprising mixing~~ [Use of] a cyst wall cysteine proteinase or a polynucleotide having a nucleic acid sequence encoding a cyst wall cysteine proteinase, ~~and a pharmaceutically acceptable carrier~~ [for the preparation of a vaccine].

Appendix A

Page A-2

Applicant(s): Salman Baig et al.

Serial No. Unknown (National Stage of PCT/US00/10672)

Filed: Herewith (Int'l. Filing Date: April 20, 2000)

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

6. (AMENDED) A vaccine comprising at least one component selected from the group consisting of (a) a polypeptide comprising a cyst wall cysteine proteinase or an immunogenic polypeptide subunit thereof and (b) a polynucleotide comprising a nucleotide sequence encoding a polypeptide comprising a cyst wall cysteine proteinase or an immunogenic polypeptide subunit thereof, and a pharmaceutically acceptable carrier.

7. The vaccine of claim 6 wherein the cyst wall cysteine proteinase is derived from *T. solium* or *T. crassiceps*.

8. (AMENDED) A method for treating an animal harboring a *Taenia* infection comprising administering to the infected animal the vaccine of claim 6 [claims 6 or 7], wherein administration of the vaccine is effective to eliminate the parasite from the animal or to prevent or delay the appearance of cysticercosis or neurocysticercosis in the animal.

9. (AMENDED) A method for protecting an animal against a *Taenia* infection comprising administering to an uninfected animal the vaccine of claim 6 [claims 6 or 7], wherein administration of the composition is effective to prevent subsequent infection of the animal by the parasite or to prevent the development of cysticercosis or neurocysticercosis in the animal after subsequent infection by the parasite.

10. (AMENDED) The method of claim 8 [claims 8 or 9] wherein the animal is a pig or a human and the *Taenia* infection is a *T. solium* infection.

11. A pharmaceutical composition comprising:

- an inhibitor molecule that inhibits the activity of a cyst wall cysteine protease, the
- inhibitor molecule comprising a peptide or peptidomimetic compound; and
- a pharmaceutically acceptable carrier.

Appendix A

Page A-3

Applicant(s): Salman Baig et al.

Serial No. Unknown (National Stage of PCT/US00/10672)

Filed: Herewith (Int'l. Filing Date: April 20, 2000)

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

12. The pharmaceutical composition of claim 11 wherein the peptide or peptidomimetic compound comprises (Xaa)_n-Yaa-Zaa-R; wherein Xaa and Zaa are each independently any amino acid; Yaa is a hydrophobic amino acid; R comprises a nucleophilic moiety; and n = 0-5.

13. The pharmaceutical composition of claim 12 wherein Yaa is leucine.

14. (AMENDED) The pharmaceutical composition of claim 12 [claims 12 or 13] wherein R comprises a chemical moiety selected from the group consisting of a carboxylic acid derivative, an amide derivative, a benzene derivative, a phenyl derivative, a chloromethylketone or derivative thereof, a fluoromethylketone or derivative thereof, an alphaketo acid or derivative thereof, a ketoamide or derivative thereof, a ketoester or derivative thereof, a vinylsulfone or derivative thereof, and a pyridyl or derivative thereof.

15. (AMENDED) The pharmaceutical composition of claim 12 [claims 12-14] wherein R comprises a fluoromethylketone (FMK).

16. The pharmaceutical composition of claim 15 wherein the inhibitor molecule is Z-Leu-Leu-Leu-FMK or Leu-Leu-Tyr-FMK.

17. (AMENDED) The pharmaceutical composition of claim 11 [16] wherein the inhibitor molecule inhibits cyst wall cysteine proteinase derived from *T. solium* or *T. crassiceps*.

18. (AMENDED) A pharmaceutical composition comprising Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK, and a pharmaceutically acceptable carrier [for use as an active pharmaceutical substance].

Appendix A

Page A-4

Applicant(s): Salman Baig et al.

Serial No. Unknown (National Stage of PCT/US00/10672)

Filed: Herewith (Int'l. Filing Date: April 20, 2000)

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

19. (AMENDED) A method for [Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK according to claim 18 for use in] treating human neurocysticercosis [comprising administering to a human subject the pharmaceutical composition of claim 18] [or porcine cysticercosis].

20. (AMENDED) A method for treating porcine cysticercosis comprising administering to a porcine subject the [The use of Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK for the preparation of a] pharmaceutical composition of claim 18.

21. (AMENDED) A method for treating human neurocysticercosis or porcine cysticercosis [comprising administering to a human or porcine subject a pharmaceutical composition comprising [The use of] an inhibitor molecule that inhibits the activity of a Taenia cysteine protease, the inhibitor molecule comprising a peptide or peptidomimetic compound, and a pharmaceutically acceptable carrier], for the treatment of human neurocysticercosis or porcine cysticercosis].

22. (AMENDED) The [method of [use of an inhibitor molecule according to] claim 21 wherein the peptide or peptidomimetic compound comprises (Xaa)_n-Yaa-Zaa-R; wherein Xaa and Zaa are each independently any amino acid; Yaa is a hydrophobic amino acid; R comprises a nucleophilic moiety; and n = 0-5.

23. (AMENDED) The [method of [use of an inhibitor molecule according to] claim 22 wherein Yaa is leucine.

24. (AMENDED) The [method of claim 22 [use of an inhibitor molecule according to claims 22 or 23] wherein R comprises a chemical moiety selected from the group consisting of a carboxylic acid derivative, an amide derivative, a benzene derivative, a phenyl derivative, a

Appendix A

Page A-5

Applicant(s): Salman Baig et al.

Serial No. Unknown (National Stage of PCT/US00/10672)

Filed: Herewith (Int'l. Filing Date: April 20, 2000)

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

chloromethylketone or derivative thereof, a fluoromethylketone or derivative thereof, an alphaketo acid or derivative thereof, a ketoamide or derivative thereof, a ketoester or derivative thereof, a vinylsulfone or derivative thereof, and a pyridyl or derivative thereof.

25. (AMENDED) The method of claim 22 [use of an inhibitor molecule according to claims 22-24] wherein R comprises a fluoromethylketone (FMK).

26. A method for inhibiting *Taenia* cyst wall cysteine protease activity comprising contacting a *Taenia* cyst wall cysteine protease with an inhibitor molecule comprising $(Xaa)_n$ -Yaa-Zaa-R; wherein Xaa and Zaa are each any amino acid; Yaa is a hydrophobic amino acid; R comprises a nucleophilic moiety; and n is 0 to about 5.

27. The method of claim 26 wherein Yaa is leucine.

28. (AMENDED) The method of claim 26 [claims 26 or 27] wherein R comprises a chemical moiety selected from the group consisting of a carboxylic acid derivative, an amide derivative, a benzene derivative, a phenyl derivative, a chloromethylketone or derivative thereof, a fluoromethylketone or derivative thereof, an alphaketo acid or derivative thereof, a ketoamide or derivative thereof, a ketoester or derivative thereof, a vinylsulfone or derivative thereof, and a pyridyl or derivative thereof.

29. (AMENDED) The method of claim 26 [28] wherein R comprises a fluoromethylketone (FMK).

30. The method of claim 29 wherein the inhibitor molecule is Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK.

Appendix A

Page A-6

Applicant(s): Salman Baig et al.

Serial No. Unknown (National Stage of PCT/US00/10672)

Filed: Herewith (Int'l. Filing Date: April 20, 2000)

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF
NEUROCYSTICERCOSIS

31. (AMENDED) The method of claim 26 [REDACTED] performed in cell free environment.
32. (AMENDED) The method of claim 26 [REDACTED] performed in cell culture, in an organ, or in a tissue.
33. (AMENDED) The method of claim 26 [REDACTED] performed in a whole animal.
34. (AMENDED) The method of claim 26 [REDACTED] wherein the *Taenia* cysteine proteinase is derived from *T. solium*.
35. A method for identifying an inhibitor of *Taenia* cysteine proteinase activity comprising:
combining a candidate inhibitor and a *Taenia* cysteine proteinase to form a mixture;
adding the proteinase substrate Z-Phe-Arg-7-amino-4-trifluoromethylcoumarin to the mixture; and
determining the extent to which the proteinase substrate is cleaved;
wherein a reduction in the extent of substrate cleavage compared to the extent of substrate cleavage in the absence of the candidate inhibitor is indicative of inhibition of *Taenia* cysteine proteinase activity.
36. The method of claim 35 comprising combining a candidate inhibitor and a cysteine protease derived from *T. solium* or *T. crassiceps* to form the mixture.
37. A computer-assisted method for identifying an inhibitor of *Taenia* cysteine proteinase activity comprising:
supplying a computer model of the structure of an inhibitor of *Taenia* cysteine proteinase activity;

Appendix A

Page A-7

Applicant(s): Salman Baig et al.

Serial No. Unknown (National Stage of PCT/US00/10672)

Filed: Herewith (Int'l. Filing Date: April 20, 2000)

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF
NEUROCYSTICERCOSIS

either (i) computationally or visually screening a structural library for candidate compounds having a structure similar to that of the inhibitor or (ii) computationally or visually designing a candidate compound having a structure similar to that of the inhibitor; and
assaying the candidate compound for the ability to inhibit *Taenia* cysteine proteinase activity.

38. A computer-assisted method for identifying an inhibitor of *Taenia* cysteine proteinase activity comprising:

solving the X-ray crystal structure of a *Taenia* cysteine proteinase to yield a computer model of the *Taenia* cysteine proteinase;

supplying a computer model of the structure of an inhibitor of *Taenia* cysteine proteinase activity;

computationally or visually docking the inhibitor structure to the proteinase crystal structure at the binding site of the inhibitor;

computationally or visually identifying intermolecular interactions between the inhibitor and the proteinase;

either (i) computationally or visually screening a structural library for candidate compounds having a structure similar to that of the inhibitor or (ii) computationally or visually designing a candidate compound having a structure similar to that of the inhibitor;

computationally or visually evaluating the intermolecular interactions between the candidate compound and the proteinase; and

assaying the candidate compound for the ability to inhibit *Taenia* cysteine proteinase activity.

39. (AMENDED) The method of **claim 37 [claims 37 or 38]** wherein the inhibitor is Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK.

Appendix A

Page A-8

Applicant(s): Salman Baig et al.

Serial No. Unknown (National Stage of PCT/US00/10672)

Filed: Herewith (Int'l. Filing Date: April 20, 2000)

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF
NEUROCYSTICERCOSIS

40. A method for treating an animal harboring a *Taenia* infection comprising administering to the infected animal a pharmaceutical composition comprising an inhibitor of a *Taenia* cysteine proteinase, wherein administration of the composition is effective to eliminate the parasite from the animal or to prevent or delay the appearance of cysticercosis or neurocysticercosis the animal.

41. A method for protecting an animal against a *Taenia* infection comprising administering to an uninfected animal a pharmaceutical composition comprising an inhibitor of a *Taenia* cysteine proteinase, wherein administration of the composition is effective to prevent subsequent infection of the animal by the parasite or to prevent the development of cysticercosis or neurocysticercosis the animal after subsequent infection by the parasite.

42. (AMENDED) The method of claim 40 [claims 40 or 41] wherein inhibitor is Z-Leu-Leu-Leu-FMK or Z-Leu-Leu- Tyr-FMK.

43. (AMENDED) The method of claim 40 [claims 40, 41 or 42] wherein the animal is a pig or a human and the *Taenia* infection is a *T. solium* infection.

44. (NEW) The method of claim 9 wherein the animal is a pig or a human and the *Taenia* infection is a *T. solium* infection.

45. (NEW) The method of claim 38 wherein the inhibitor is Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK.

46. (NEW) The method of claim 41 wherein inhibitor is Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK.

Appendix A

Page A-9

Applicant(s): Salman Baig et al.

Serial No. Unknown (National Stage of PCT/US00/10672)

Filed: Herewith (Int'l. Filing Date: April 20, 2000)

For: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF
NEUROCYSTICERCOSIS

47. (NEW) The method of claim 41 wherein the animal is a pig or a human and the *Taenia* infection is a *T. solium* infection.

48. (NEW) The method of claim 42 wherein the animal is a pig or a human and the *Taenia* infection is a *T. solium* infection.

**CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND
TREATMENT OF NEUROCYSTICERCOSIS**

5 This application claims the benefit of U.S. Provisional Application Serial No. 60/130,338, filed April 21, 1999, which is incorporated herein by reference in its entirety.

Statement of Government Rights

10 This invention was made with government support under a grant from the US/Mexico Foundation, Grant No. 42-H-94, and the N.I.H. Molecular Parasitology Training Grant For Cell Biology of Parasites and Vectors, Grant No. AIO7322. The U.S. Government may have certain rights in this invention.

Background

15 Neurocysticercosis is one of the most common parasitic disease of the human central nervous system and the leading cause of epileptiform seizures in many parts of the Third World. Presently, no completely efficacious treatments nor vaccines for neurocystercercosis exist (Garcia et al., *J. Infec. Dis.* 175:486-
20 489 (1997)). Caused by cysts of the pork tapeworm, *Taenia solium*, the disease is estimated to affect 50 million people worldwide, although some epidemiological studies suggest that this figure may be underestimated. It is the most likely reason that epilepsy is twice as common in developing as opposed to developed countries of the world. Some symptoms associated with
25 neurocysticercosis include seizures, headaches, visual problems, confusion, and psychosis. However, there is a prepatant asymptomatic period which can last for 10 years or more, and usually approximates the life expectancy of the tissue cysts. Evidence suggests that tissue cysts suppress host inflammatory responses to complete their life cycles, whereas clinical symptoms emerge during the

interval between cyst degeneration and death (Evans et al., Emerg. Infect. Dis. 3:403-405 (1997)).

With the increase in immigration, cysticercosis is an emerging infectious disease in the United States. The number of neurocystercosis cases in the U.S. is estimated to be greater than 1000 per year, and this number appears to be on the rise. For example, 10% of neurology admissions at Los Angeles County USC Medical Center in the past three years were for neurocystercosis, which amounted to about 120 cases/year. Surely, the underlying rate of infection is higher. Locally acquired cases have also been documented in Denver, San Diego, New York City, Chicago, and other unexpected locations. For example, 1.3% of an orthodox New York Jewish community were found to test positive for neurocystercosis. Although orthodox Jews would not be expected to be at risk for infection because they shun pork, they were reported to have been infected through contamination from their immigrant Latino domestic helpers who were *Taeniasis* (tapeworm carriers). Four million people worldwide are reportedly infected with *Taeniasis*, harboring the adult tapeworm stage of *T. solium*. For every one of these victims, it is estimated that 10 or more people are infected with the cyst stage, i.e., are cysticercotic. In areas of high endemicity, almost anyone is at risk for neurocystercosis, since the eggs can pervade the environment.

Humans are the only definitive host for the adult pork tapeworm, which remains in the intestine attached with its scolex. Humans acquire neurocystercosis when they become accidental *intermediate* hosts to the cyst form of the parasite, commonly acquiring eggs through human fecal contamination or by direct contact with a tapeworm carrier. Pigs are the normal intermediate hosts. In humans, the cysts tend to localize more in the central nervous system, hence the prevalence of neurocystercosis.

The International Task Force for Disease Eradication has identified neurocystercosis as a target for eradication efforts since *T. solium* has only two hosts: humans and pigs. Although good porcine husbandry has met with

some success in interrupting transmission of *T. solium* in western Europe and the U.S., these approaches are doomed to failure in developing countries since poor farmers often prefer to sell infected pigs in an informal setting, rather than turning them over for possible condemnation. Some people are reported to actually prefer cysticercotic ("measly") pork to eat.

To date, no vaccines have been developed for any human parasites, including *T. solium*. Recent research has indicated that mass chemotherapy approaches are more promising and practical for the purposes of neurocystercercosis eradication. For example, an initial mass treatment with Praziquantel(TM) in Ecuador caused porcine cysticercosis to fall from 11.4 to 2.6%. Present treatment for active parenchymal neurocystercercosis commonly employs the antiparasitic compounds Praziquantel, Albendazole(TM), or a combination of both. A slight effect toward cyst resolution has been noted in patients who were treated with Praziquantel followed by subsequent treatment with Albendazole. However, some patients presented with worsening inflammatory reactions around the cysts with symptoms including nausea, vomiting, and headache; some patients developed cerebral edema. While some reports have shown that these drugs have demonstrated control of seizures in neurocystercercosis patients, it is not clear if these findings were a result of therapy or a result of patient selection bias. Consequently, the risks of developing acute inflammation, cerebral edema, and possibly chronic hydrocephalus associated with these agents is of significant concern. Although some studies suggest that Albendazole may be more effective than Praziquantel, it has also been linked to hematopoietic and hepatic dysfunction and may be teratogenic/mutagenic. Carpio and colleagues conducted the largest single randomized and blinded trial treatment of neurocystercercosis with Albendazole and Praziquantel, reported to date, and conclude that any benefits appear to be related to the natural course of the infection rather than any specific effect of drug treatment (Arch. Int. Med., 155: 1982-1988 (1995)). There appear to be no antiparasitic drugs that have been developed exclusively for treatment of

neurocystercercosis. Clearly, there is a need for a more efficacious and specific anti-neurocystercercosis drug, possibly to be optimized to work as an adjunct to existing therapies.

5

Summary of the Invention

A cyst wall cysteine proteinase has been discovered in the parasite *T. solium* that appears to play a significant biological role in the organism. The invention is thus directed to the *T. solium* cyst wall cysteine proteinase and a polynucleotide having a nucleotide sequence that encodes the cysteine proteinase (and its complement). The invention is further directed to a vaccine that includes either or both of a cyst wall cysteine proteinase (or an immunogenic polypeptide subunit thereof) or a polynucleotide having a nucleotide sequence encoding a cyst wall cysteine proteinase (or an immunogenic polypeptide subunit thereof). The cyst wall cysteine proteinase used in the vaccine is preferably derived from *T. solium* or *T. crassiceps*. Also included in the invention are methods of preparing and using the vaccine.

Inhibitors of the activity of cyst wall cysteine proteinase have also been discovered. The invention is thus also directed to a pharmaceutical composition that includes an inhibitor molecule that inhibits the activity of a cyst wall cysteine protease, together with a pharmaceutically acceptable carrier. The inhibitor molecule is preferably a peptide or peptidomimetic compound, and preferably inhibits cyst wall cysteine proteinase derived from *T. solium* or *T. crassiceps*. In a preferred embodiment, the inhibitor peptide or peptidomimetic compound includes (Xaa)_n-Yaa-Zaa-R; wherein Xaa and Zaa are each independently any amino acid; Yaa is a hydrophobic amino acid; R comprises a nucleophilic moiety; and n = 0-5. More preferably, the inhibitor peptide or peptidomimetic compound contains a blocking group at the N-terminus. The pharmaceutical composition is useful for treating human neurocystercercosis or porcine cysticercosis. Methods of preparing and using the inhibitory compound

and the pharmaceutical composition are also included in the invention. For example, the invention includes a method for inhibiting cyst wall cysteine proteinase activity that involves contacting a cyst wall cysteine proteinase with an inhibitor molecule. The method of inhibiting cyst wall cysteine proteinase can be performed in cell free environment, in cell culture, in an organ, in a tissue, or in a whole animal, such as a human or a pig.

Also included in the invention is a method for identifying an inhibitor of *Taenia* cysteine proteinase activity. A candidate inhibitor is combined with a *Taenia* cysteine proteinase to form a mixture. The proteinase substrate Z-Phe-Arg-7-amino-4-trifluoromethylcoumarin is then added to the mixture, and the extent to which the proteinase substrate is cleaved is determined. It should be understood that the components of the mixture can be added simultaneously or in any desired order. A reduction the extent of substrate cleavage compared to the extent of substrate cleavage in the absence of the candidate inhibitor is indicative of inhibition of *Taenia* cysteine proteinase activity.

The invention also includes a computer-assisted method for identifying an inhibitor of *Taenia* cysteine proteinase activity. In one embodiment, a computer model of the structure of an inhibitor of *Taenia* cysteine proteinase activity is supplied, then a structural library computationally or visually screened for candidate compounds having a structure similar to that of the inhibitor. Alternatively, a candidate compound having a structure similar to that of the inhibitor can be designed. Optionally, the candidate compound is for the ability to inhibit *Taenia* cysteine proteinase activity. In another embodiment of the computer-assisted method of drug design, the X-ray crystal structure of a *Taenia* cysteine proteinase is solved to yield a computer model of the *Taenia* cysteine proteinase, and a model of an inhibitor molecule is computationally or visually docked to the proteinase crystal structure at the binding site of the inhibitor. Intermolecular interactions between the inhibitor and the proteinase are then computationally or visually identified, and either a structural library is computationally or visually screened for candidate compounds having a structure

similar to that of the inhibitor or a candidate compound is computationally designed. Then, the intermolecular interactions between the candidate compound and the proteinase are computationally or visually evaluated, and the candidate compound is optionally assayed for the ability to inhibit *Taenia* cysteine proteinase activity.

Brief Description of the Figures

Figure 1 shows (a) 30 amino acid N-terminal sequence of *T. crassiceps* cysteine protease (SEQ ID NO:1); and (b) alignment of catalytic cysteine residue (*) with other cysteine proteases.

Figure 2 is a comparison of FPLC anion-exchange chromatography of active fractions from ACA 54 gel-filtration separations of a) *T. crassiceps* (Example I) and b) *T. solium* (Example II).

Figure 3 shows substrate specificity of the purified a) *T. crassiceps* (Example I) and b) *T. solium* (Example II) cysteine proteases assessed through cleavage of a panel of peptide substrates with different substitutions at P1-P3.

Figure 4 is Western blot showing degradation of IgG heavy chain by cysteine proteases of a) *T. crassiceps* (Example I) and b) *T. solium* (Example II).

Figure 5 shows stimulation of splenocytes removed from *T. crassiceps* infected mice by the purified *T. solium* enzyme.

Figure 6 shows inhibition of *T. solium* cysteine protease's cleavage of its primary substrate, Z-Phe-Arg-AFC, by various inhibitors.

Figure 7 shows scanning electron microscopy (SEM) of surface of cysts removed from mice treated with *Taenia* cysteine protease inhibitor and control mice. Magnifications are identical for both sides in each panel. Panel A (4000x): SEM shows a vigorous host immune response on cysts treated with Z-LLY-FMK. No immune cells were seen on cysts removed from untreated mice. Panel B (25,000x): SEM results show fibroblasts, holes in the tegument, and sloughed microtriches in cysts from treated mice. Microtriches in untreated cysts can be seen to be visibly longer. Visible anchors are also noted for the

untreated cyst group whereas these anchors were not observed in microtriches from treated cysts.

Figure 8 shows *in vitro* cytotoxicity analysis of Z-LLL-FMK and Z-LLY-FMK inhibitors.

5

Detailed Description of the Invention

A cysteine proteinase that is capable of degrading host proteins, including human IgG, has been discovered in the cyst wall of the human parasite *T. solium*. *Taenia* cyst wall cysteine proteinases are significant to the life cycle of the parasite, thereby presenting important chemotherapeutic targets. Cyst wall cysteine proteinases of *T. solium* and *T. crassiceps* (the mouse parasite) are biochemically similar, and inhibitor molecules have been discovered that specifically inhibit the activity of both these enzymes. Inhibitors of *T. solium* and *T. crassiceps* cysteine proteinase are useful to treat or prevent infection of animal hosts, including humans.

Accordingly, the present invention is directed to a proteinase, preferably an isolated cyst wall cysteine proteinase, derived from *Taenia solium*, which proteinase is capable of cleaving the substrate Z-Phe-Arg-7-amino-4-trifluoromethylcoumarin (Z-Phe-Arg-AFC; where Z is the blocking group $\text{CH}_2\text{-CH}_2\text{-O-(C=O)-}$ which is also known as "CBZ"). Z-Phe-Arg-AFC is commercially available from Enzyme Systems Products, Livermore CA, 94550. A protein, such as a proteinase, that is "derived from" *T. solium*, as that term is used herein, is to be broadly understood to include not only a protein that is physically isolated from *T. solium*, but also a protein having the same activity and, optionally, the same amino acid sequence as a protein thus isolated which is produced or obtained in any other manner, such as by means of chemical synthesis or recombinant DNA technology. Likewise, the term "*T. solium* protein" is to be understood to include a protein isolated from *T. solium*, together with synthetic or recombinant functional equivalents thereof. The terms

"proteinase" and "protease" are used interchangeably herein, and mean an enzyme capable of cleaving a protein at a bond, typically a peptide bond, that joins together two amino acids.

- Preferably, the proteinase of the invention has one or more of the following characteristics:

- (1) it is capable of rapidly and completely degrading the heavy and light chains of human IgG (for example, a nonlimiting amount of cysteine proteinase can completely degrade 1 ug of human IgG within about 120 minutes, preferably within about 45 minutes, as detected using a Western blot);
- (2) it has a molecular weight of about 43 kD to about 65 kD;
- (3) optimal proteinase activity occurs at acidic pH, about pH 4.8;
- (4) it is thiol-activated; that is, it requires exogenous thiol groups, such as L-cysteine, for activity;
- (5) the ratio K_{cat}/K_m using Z-Phe-Arg-AFC as a substrate is essentially at the diffusion control limit (preferably greater than about $10^8 \text{ M}^{-1}\text{s}^{-1}$)
- (6) it has only one substrate binding site;
- (7) it obeys Michaelis-Menten kinetics; that is, it is noncooperative;
- (8) it is inhibited by cysteine proteinase inhibitors, but is not inhibited to the same extent by inhibitors of other types of proteases; and

- The proteolytic activity of the newly identified proteinase of the human parasite, *T. solium*, appears to be essentially undistinguishable from the demonstrated activity of the analogous protein of the mouse parasite, *T. crassiceps* (White et al., *J. Mol. Biochem. Parasitol.*, 85:243-253 (1997)).

- The invention is further directed to an isolated nucleic acid molecule comprising a nucleic acid sequence that encodes a proteinase of the invention. The complement of such a nucleic acid molecule is also encompassed by the invention.

- An "isolated" biomolecule, such as a polypeptide or polynucleotide, is a biomolecule that has been either removed from its natural environment, produced using recombinant techniques, or chemically or enzymatically

synthesized. Preferably, a biomolecule of this invention is purified, i.e., essentially free from any other biomolecules and associated cellular products or other impurities. The terms "polypeptide" or "polypeptide subunit," as used herein, refer to a polymer of amino acids and do not connote a specific length of a polymer of amino acids. Thus, for example, the terms peptide, oligopeptide, protein, and enzyme are included within the definition of polypeptide, whether produced using recombinant techniques, chemical or enzymatic synthesis, or naturally occurring. This term also includes polypeptides that have been modified or derivatized, such as by glycosylation, acetylation, phosphorylation, and the like.

The present invention is further directed to an inhibitor of a cyst wall cysteine proteinase. The inhibitor molecule inhibits the activity of a cyst wall cysteine proteinase, preferably a *Taenia* cyst wall cysteine protease, more preferably a *T. solium* or *T. crassiceps* cyst wall cysteine proteinase. An inhibitor molecule is preferably a peptide (i.e., a molecule that contains two or amino acids or derivatized amino acids linked by a peptide bond) or a peptidomimetic compound, although it can contain one amino acid or derivatized amino acid. A "peptidomimetic" compound is a compound that functionally and/or structurally mimics a peptide, but that lacks one or more of the peptide bonds that characterize the peptide. Peptidomimetic compounds therefore not typically do not serve as substrate for proteases and are likely to be active *in vivo* for a longer period of time as compared to the analogous peptides. Unless otherwise specified, the term "peptide" when used herein in the context of an inhibitor peptide or inhibitor molecule includes peptides and peptidomimetic compounds.

In one preferred embodiment, the inhibitor molecule contains, covalently linked to its C-terminus, a nucleophilic chemical moiety. The nature of the nucleophile is not limiting and can include a carboxylic acid derivative, an amide derivative, a benzene ring derivative, a phenyl derivative, a pyridyl derivative, and the like. Suitable nucleophilic moieties include but are not limited to a

chloromethylketone (CMK), a fluoromethylketone (FMK), an alphaketo acid (AK), a ketoamide (KA), a ketoester (KE), a vinylsulfone (VS), a pyridyl group (Pyr), or any combination thereof. A preferred nucleophilic moiety is a fluoromethylketone (FMK). Preferably the nucleophilic moiety is also

- 5 hydrophobic, thereby assisting the inhibitor molecule in crossing the blood-brain barrier. Additionally or alternatively, a preferred inhibitor peptide contains, at the P2 position (P2 being the second amino acid position, counted from the C-terminus), a hydrophobic amino acid, preferably an aliphatic hydrophobic amino acid, more preferably a leucine. In a preferred embodiment, a tyrosine or leucine
- 10 is at the P1 position of the peptide; and/or leucine is at the P3 position, although it should be understood that any amino acids are well tolerated at the P1 and P3 positions. Particularly preferred inhibitor peptides include Xaa_n-Leu-Tyr-FMK and Xaa_n-Leu-Leu-FMK where n = 0-5; more preferably, n = 0-2, for example Z-Leu-Leu-Tyr-Fluoromethylketone (Z-LLY-FMK) and Z-Leu-Leu-Leu-
- 15 Fluoromethylketone (Z-LLL-FMK), commercially available from Enzyme System Products, Livermore, CA. Other preferred inhibitor peptides include (Ph)₂CHCO-Leu-Phe-CO-NHCH₂-2-Pyridyl (Ph₂-LF-KP); Z-Leu-Phe-CONH-Ethyl (Z-LF-KE); Z-Leu-Phe-CONH-4-Morphophinyl (Z-LF-KM); Z-Leu-Phe-CH₂-CH(OH)-C₆H₄-(4-OPh) (Z-LF-BPh); Boc-Leu-Phe-Vinylsulfone-Phenyl
- 20 (B-LF-VS); and Z-Leu-Leu-Leu-Vinylsulfone-Phenyl (Z-LLL-VS).

- The N-terminus of an inhibitor peptide or peptidomimetic of the invention is preferably derivatized. For example, it can be covalently linked to a blocking group, such as "Z" (CH₃-CH₂-O-(C=O)- which is also known as "CBZ") or "Boc" (t-butoxy). N-terminal blocking groups are well known in the
- 25 art of synthetic organic chemistry and peptide synthesis.

- The invention further provides methods for treating or preventing infection of an animal by *Taenia*. Preferably, the animal is a cow, a dog, a pig or a human. More preferably, the animal is a pig or a human and the method is effective for treating and preventing a *T. solium* infection. In one aspect, a
- 30 pharmaceutical composition containing an inhibitor molecule of the invention is

administered therapeutically to an animal harboring a *Taenia* infection. In one embodiment of the therapeutic method, administration of the inhibitor molecule is effective to eliminate the parasite from the animal; in another embodiment, administration of the inhibitor molecule is effective to prevent or delay the appearance of cysticercosis or neurocysticercosis in the animal. In another aspect, a pharmaceutical composition containing an inhibitor molecule of the invention is administered prophylactically to an animal in advance of infection by the parasite. In one embodiment of the prophylactic method, administration of the inhibitor molecule is effective to prevent subsequent infection of the animal by the parasite. In another embodiment of the prophylactic method, administration of the inhibitor molecule is effective to prevent or delay the development of cysticercosis or neurocysticercosis in the animal after subsequent infection by the parasite. In yet another embodiment of the prophylactic method, administration of the inhibitor molecule is effective to prevent the death of the animal after subsequent infection by the parasite.

Both veterinary and human applications are contemplated, as the inhibitor molecule is expected to inhibit the activity of cyst wall cysteine protease derived from various species of *Taenia*, including *T. solium* and *T. crassiceps*. Veterinary applications include prophylactic and therapeutic treatment of domestic animals, which may also control the spread of infection into human populations. Preferably, the method of the invention comprises administration to a pig or human that is infected with *Taenia* or at risk of *Taenia* infection. In humans, the inhibitor is preferably introduced into the bloodstream, for example by intravenous injection or by ingestion. In pigs, the inhibitor is preferably administered in feed as a prophylactic or therapeutic agent against porcine cysticercosis, with the ultimate goal of interrupting the life cycle of *Taenia* at the porcine stage. Such a strategy could conceivably lead to the ultimate eradication of human neurocysticercosis.

Example V describes the use of inhibitor peptides of the invention to treat cysticercosis in the murine model of *Taenia* infection. These studies show

some of the highest reported protection levels in mice challenged with cysticerci, and the inhibitors were highly efficacious at low concentration levels.

Moreover, mice which were treated with the inhibitor displayed no apparent toxicity effects upon visual examination. The results of these mouse studies

- 5 strongly suggest that the inhibitor peptides are specific for a cyst wall cysteine protease which is present only in the parasite *Taenia*. In other words, the inhibitor peptide is not expected to significantly interfere with the activity of proteolytic enzymes in a mammalian host, such as a pig or a human. Accordingly, treatment of an infected animal with the inhibitor molecule of the
- 10 invention is expected to be more specific than present treatments for neurocysticercosis, which include administration of Praziquantel and Albendazole. Both of these drugs were discovered by drug screening approaches and not by an approach directed toward inhibition of a particular *Taenia* enzyme activity, as in the present invention. Both drugs are not very
- 15 effective and are accompanied by significant side effects in hosts.

The inhibitor molecules of the invention are readily formulated as pharmaceutical compositions for veterinary or human use. The pharmaceutical composition optionally includes excipients or diluents that are pharmaceutically acceptable as carriers and compatible with the inhibitor molecule. The term

20 "pharmaceutically acceptable carrier" refers to a carrier(s) that is "acceptable" in the sense of being compatible with the other ingredients of a composition and not deleterious to the recipient thereof. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the pharmaceutical composition may contain minor

25 amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, salts, and/or adjuvants which enhance the effectiveness of the immune-stimulating composition. In a preferred embodiment, the pharmaceutical composition is formulated so as to allow or enhance transmission of the inhibitor molecule across the blood/brain barrier in humans. Methods of

making and using such pharmaceutical compositions are also included in the invention.

The inhibitor molecules can be formulated in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active compound into association with a carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing the active compound into association with a liquid carrier, a finely divided solid carrier, or both, and then, if necessary, shaping the product into the desired formulations.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as tablets, troches, capsules, lozenges, wafers, or cachets, each containing a predetermined amount of the inhibitor as a powder or granules, as liposomes containing the inhibitor, or as a solution or suspension in an aqueous liquor or non-aqueous liquid such as a syrup, an elixir, an emulsion or a draught. Such compositions and preparations should contain at least about 0.1% active compound. The percentage of the compositions and preparations may be varied and may conveniently be between about 1% to about 60% of the weight of a given unit dosage form.

Tablets, troches, pills, capsules, and the like may also contain one or more of the following: a binder such as gum tragacanth, acacia, corn starch or gelatin; an excipient such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; a sweetening agent such as sucrose, fructose, lactose or aspartame; and a natural or artificial flavoring agent. When the unit dosage form is a capsule, it may further contain a liquid carrier, such as a vegetable oil or a polyethylene glycol. Various other materials may be present as coatings or to otherwise modify the physical form of the solid unit dosage form. For instance, tablets, pills, or capsules may be coated with gelatin, wax, shellac, or sugar and the like. A syrup or elixir may contain one or more of a sweetening agent, a preservative such as methyl- or propylparaben, an agent to retard crystallization

of the sugar, an agent to increase the solubility of any other ingredient, such as a polyhydric alcohol, for example glycerol or sorbitol, a dye, and flavoring agent. The material used in preparing any unit dosage form is substantially nontoxic in the amounts employed. The inhibitor may be incorporated into sustained-release
5 preparations and devices.

The inhibitors of the invention can be incorporated directly into the food of the animal's diet, as an additive, supplement, or the like. Thus, the invention further provides a food product containing an inhibitor of the invention. Any food is suitable for this purpose, although processed foods already in use as
10 sources of nutritional supplementation or fortification, such as animal feeds, breads, cereals, and the like, may be more convenient to use for this purpose.

Formulations suitable for parenteral administration conveniently comprise a sterile aqueous preparation of the inhibitor, or dispersions of sterile powders comprising the inhibitor, which are preferably isotonic with the blood
15 of the recipient. Isotonic agents that can be included in the liquid preparation include sugars, buffers, and sodium chloride. Solutions of the inhibitor can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions of the inhibitor can be prepared in water, ethanol, a polyol (such as glycerol, propylene glycol, liquid polyethylene glycols, and the like), vegetable oils,
20 glycerol esters, and mixtures thereof. The ultimate dosage form is sterile, fluid and stable under the conditions of manufacture and storage. The necessary fluidity can be achieved, for example, by using liposomes, by employing the appropriate particle size in the case of dispersions, or by using surfactants. Sterilization of a liquid preparation can be achieved by any convenient method
25 that preserves the activity of the inhibitor, preferably by filter sterilization. Preferred methods for preparing powders include vacuum drying and freeze drying of the sterile injectible solutions. Subsequent microbial contamination can be prevented using various antimicrobial agents, for example, antibacterial, antiviral and antifungal agents including parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. Delivery of the inhibitors over a prolonged period
30

can be achieved by including agents for delaying, for example, aluminum monostearate and gelatin.

Nasal spray formulations comprise purified aqueous solutions of the inhibitor with preservative agents and isotonic agents. Such formulations are preferably adjusted to a pH and isotonic state compatible with the nasal mucous membranes.

Formulations for rectal or vaginal administration may be presented as a suppository with a suitable carrier such as cocoa butter, or hydrogenated fats or hydrogenated fatty carboxylic acids.

Ophthalmic formulations are prepared by a similar method to the nasal spray, except that the pH and isotonic factors are preferably adjusted to match that of the eye.

Topical formulations comprise the inhibitor dissolved or suspended in one or more media such as mineral oil, petroleum, polyhydroxy alcohols or other bases used for topical pharmaceutical formulations. The compound of the invention is particularly suited to incorporation in a cosmetic lotion, crème, or sunscreen for use on the skin.

Useful dosages of the inhibitor molecule of the invention can be determined by comparing their *in vitro* activity and the *in vivo* activity in animals models. Methods for extrapolation of effective dosages of drugs in mice, and other animals, to humans are known in the art; for example, see U.S. Pat. No. 4,938,949, which is incorporated herein by reference, in its entirety.

It is expected that the inhibitor molecule of the invention will be superior to Praziquantel and Albendazole for treatment of neurocysticercosis as a result of its high efficacy and apparent specificity, which may lead to the effective use of lower drug dosages compared to those used for Praziquantel and Albendazole. In some applications, a treatment protocol may include administration of a combination of Praziquantel, Albendazole and/or an inhibitor molecule of the invention.

It should be understood that an inhibitor molecule of the invention may be effective to treat other parasitic infections, if the active site of the cyst wall cysteine proteinase is conserved among other metabolically important enzymes in other parasites.

- 5 The invention additionally provides a method for identifying an inhibitor of cyst wall cysteine proteinase activity. A candidate inhibitor is combined with a cyst wall cysteine proteinase derived from *Taenia*, preferably *T. solium* or *T. crassiceps*, to form a mixture. A detectable substrate for the proteinase, preferably the fluorogenic molecule Z-Phe-Arg-AFC, is then added to the mixture. After a period of time and under conditions sufficient to allow for
10 cleavage, as described more fully hereinbelow, it is determined whether or not the substrate is cleaved. The absence of cleavage, i.e., recovery of intact substrate, identifies a promising inhibitor of cyst wall cysteine proteinase activity.

- 15 The three-dimensional conformations of the inhibitor peptides Z-LLY-FMK, Z-LLL-FMK, Ph2-LF-KP or any of the other inhibitor peptides described herein are amenable to computer-enhanced display and manipulation. These structures can be computationally constructed from readily accessible peptide libraries. It is within the scope of the invention to identify conformational
20 analogues to Z-LLY-FMK or Z-LLL-FMK, for example, having different compositions, using computer-aided rational drug design techniques in common use in the field of computational biology. Computer-generated structural libraries of peptides and other small molecules are readily available, and can be computer-screened for conformational similarity to Z-LLY-FMK, or Z-LLL-FMK.
25 Optionally, the X-ray crystal structure of a *Taenia* cysteine proteinase can be determined using methods well known in the art, and the coordinates thereof can be displayed on a computer. The active site can be identified, and potential inhibitors can be identified by putative binding to the active site via computer modeling techniques. Compounds thus identified as structurally similar to

Z-LLY-FMK or Z-LLL-FMK, or other compounds identified as potentially binding to the active site of the proteinase, are then subjected to *in vitro* testing according to the invention to determine whether they inhibit cyst wall cysteine proteinase activity as described below. The invention is understood to further include inhibitor molecules identified according to the methods of the invention, in addition to the methods of identifying them.

The invention further provides a vaccine that is effective to treat or prevent infection of an animal by a cyst-forming parasite, such as *Taenia*. The vaccine can be a polypeptide vaccine or a polynucleotide vaccine, and can include one or more immunogenic components. The polynucleotide vaccine contains at least one polynucleotide having a nucleotide coding region that encodes a cyst wall cysteine proteinase or an immunogenic polypeptide subunit thereof. Analogously, a polypeptide vaccine contains at least one cyst wall cysteine proteinase or immunogenic polypeptide subunit thereof. An immunogenic polypeptide subunit of a cyst wall cysteine protease is one that elicits in an animal host an antibody-mediated immune response (i.e., a B cell response or humoral immunity), a cell-mediated immune response (i.e., a T cell response), or a combination thereof. The immunogenicity of a cysteine protease subunit can be evaluated, for example, using the assays set forth in the Examples. Optionally, the polypeptide or polynucleotide vaccine includes additional compounds that stimulate the host's immune system, as further described below. The invention should be understood as also including methods of making and using the polynucleotide and polypeptide vaccines.

The polynucleotide vaccine can contain DNA, RNA, a modified nucleic acid, or any combination thereof. Preferably, the polynucleotide vaccine comprises one or more cloning or expression vectors; more preferably, the vaccine comprises one or more expression vectors each capable of autonomous expression of a nucleotide coding region in a mammalian cell to produce at least one immunogenic polypeptide or cytokine, as further described below. An expression vector preferably includes a eukaryotic promoter sequence, more

preferably the nucleotide sequence of a strong eukaryotic promoter, operably linked to one or more coding regions. A promoter is a DNA fragment that acts as a regulatory signal and binds RNA polymerase in a cell to initiate transcription of a downstream (3' direction) coding sequence; transcription is the formation of an RNA chain in accordance with the genetic information contained in the DNA. A promoter is "operably linked" to a nucleic acid sequence if it is does, or can be used to, control or regulate transcription of that nucleic acid sequence. The invention is not limited by the use of any particular eukaryotic promoter, and a wide variety are known; preferably, however, the expression vector contains a CMV or RSV promoter. The promoter can be, but need not be, heterologous with respect to the host cell. The promoter used is preferably a constitutive promoter.

A vector useful in the polynucleotide vaccine of the present invention can be circular or linear, single-stranded or double stranded and can be a plasmid, cosmid, or episome but is preferably a plasmid. In a preferred embodiment, each nucleotide coding region (whether it encodes an immunogenic polypeptide or a cytokine) is on a separate vector; however, it is to be understood that one or more coding regions can be present on a single vector, and these coding regions can be under the control of a single or multiple promoters.

There are numerous plasmids known to those of ordinary skill in the art useful for the production of polynucleotide vaccines. Preferred embodiments of the polynucleotide vaccine of the invention employ constructs using the plasmids VR1012 (Vical Inc., San Diego CA), pCMVI.UBF3/2 (S. Johnston, University of Texas) or pcDNA3.1 (InVitrogen Corporation, Carlsbad, CA) as the vector. Plasmids VR1012 and pCMVI.UBF3/2 are particularly preferred. In addition, the vector construct can contain immunostimulatory sequences (ISS), such as unmethylated dCpG motifs, that stimulate the animal's immune system. Other possible additions to the polynucleotide vaccine constructs include nucleotide sequences encoding cytokines, such as granulocyte macrophage

colony stimulating factor (GM-CSF), interleukin-12 (IL-12) and co-stimulatory molecules such B7-1, B7-2, CD40. The cytokines can be used in various combinations to fine-tune the response of the animal's immune system, including both antibody and cytotoxic T lymphocyte responses, to bring out the specific level of response needed to control or eliminate the *Taenia* infection. The polynucleotide vaccine can also encode a fusion product containing the immunogenic polypeptide and a molecule, such as CTLA-4, that directs the fusion product to antigen-presenting cells inside the host. Plasmid DNA can also be delivered using attenuated bacteria as delivery system, a method that is suitable for DNA vaccines that are administered orally. Bacteria are transformed with an independently replicating plasmid, which becomes released into the host cell cytoplasm following the death of the attenuated bacterium in the host cell. An alternative approach to delivering the polynucleotide to an animal involves the use of a viral or bacterial vector. Examples of suitable viral vectors include adenovirus, polio virus, pox viruses such as vaccinia, canary pox, and fowl pox, herpes viruses, including catfish herpes virus, adenovirus-associated vector, and retroviruses. Exemplary bacterial vectors include attenuated forms of *Salmonella*, *Shigella*, *Edwardsiella ictaluri*, *Yersinia ruckerii*, and *Listeria monocytogenes*.

The polynucleotide vaccine of the invention can be administered to the animal using any convenient method, such as intramuscular injection, topical or transdermal application to the animal's skin, or use of a gene gun, wherein particles coated with the polynucleotide vaccine are shot into the animal's skin. The amount of polynucleotide administered to the animal is affected by the nature, size and disease state of the animal as well as the delivery method; for example, typically less DNA is required for gene gun administration than for intramuscular injection. Further, if a polynucleotide encoding a cytokine is co-delivered with nucleotide coding regions encoding the immunogenic polypeptide, the amount of polynucleotide encoding the immunogenic polypeptide from in the vaccine is optionally reduced.

Hundreds of publications have now reported the efficacy of DNA vaccines in small and large animal models of infectious diseases, cancer and autoimmune diseases (J. Donnelly et al., Rev. Immunol. 15:617 (1997)). Vaccine dosages for humans can be readily extended from the murine models by one skilled in the art of genetic immunization, and a substantial literature on genetic immunization of humans is now available to the skilled practitioner. For example, Wang et al. (Science 282:476-480 (1998)) vaccinated humans with plasmid DNA encoding a malaria protein, and the same group has developed a plan for manufacturing and testing the efficacy of a multigene *Plasmodium falciparum* liver-stage DNA vaccine in humans (Hoffman et al., Immunol. Cell Biol. 75:376 (1997)). In general, the polynucleotide vaccine of the invention is administered in dosages that contain the smallest amount of polynucleotide necessary for effective immunization. It is typically administered to human subjects in dosages containing about 20 µg to about 2500 µg plasmid DNA; in some instances 500 µg or more of plasmid DNA may be indicated. Typically the vaccine is administered in two or more injections at time intervals, for example at four week intervals.

The polypeptide vaccine of the invention, in addition to containing cyst wall cysteine proteinase or an immunogenic polypeptide subunit thereof, can include an adjuvant in order to further stimulate the animal's immune system. Adjuvants suitable for human and veterinary use are well known in the medical arts. Like the polynucleotide vaccine, the polypeptide vaccine can be administered to the animal using any convenient method, such as intramuscular or intraperitoneal injection, topical administration, oral or intranasal administration, inhalation, perfusion and the like, as described above for the administration of the inhibitor molecule. The amount of polypeptide administered to the animal is affected by the nature, size and disease state of the animal, as well as by the delivery method. Useful dosages of the polypeptide vaccine for humans can be readily determined by evaluating its activity *in vivo* activity in mice.

Accordingly, the invention further provides a method for vaccinating an animal against infection by *Taenia*. Preferably, the animal is a cow, a dog, a pig or a human. More preferably, the animal is a pig or a human and the method is effective for treating and preventing a *T. solium* infection. In one aspect, the vaccine is administered therapeutically to an animal harboring a *Taenia* infection. In one embodiment of the therapeutic method, administration of the vaccine is effective to eliminate the parasite from the animal; in another embodiment, administration of the vaccine is effective to prevent or delay the appearance of cysticercosis or neurocysticercosis in the animal. In another aspect, the vaccine is administered prophylactically to an animal in advance of infection by the parasite. In one embodiment of the prophylactic method, administration of the vaccine is effective to prevent subsequent infection of the animal by the parasite. In another embodiment of the prophylactic method, administration of the vaccine is effective to prevent or delay the development of cysticercosis or neurocysticercosis in the animal after subsequent infection by the parasite. In yet another embodiment of the prophylactic method, administration of the vaccine is effective to prevent the death of the animal after subsequent infection by the parasite. Both veterinary and human applications are contemplated, as the vaccine is expected to be effective against various species of *Taenia*, including *T. solium* and *T. crassiceps*. Veterinary applications include prophylactic and therapeutic treatment of domestic animals, which may also control the spread of infection into human populations. Preferably, the method of the invention comprises administration of the vaccine to a pig or human that is infected with *Taenia* or at risk of *Taenia* infection. In pigs, the polynucleotide or polypeptide vaccine is preferably administered in feed.

The invention further contemplates administration to an animal of different vaccines and/or inhibitor molecules in a serial protocol. For example, a plasmid-based DNA vaccine as described herein can be administered to a animal to "prime" the immune system, followed by the one or more administrations of a

polypeptide vaccine, a viral vaccine (e.g., vaccinia vector carrying the genes that encode the immunogenic polypeptide and, optionally, cytokines) and/or a pharmaceutical composition comprising an inhibitor of cyst wall cysteine protease activity as identified according to the invention. The order of administration of the different types of compositions, and the nature of the compositions administered in any given dose (e.g., polypeptide vaccine, plasmid vaccine, viral vector vaccine, inhibitor molecule) can be readily determined by one of skill in the art to invoke the most effective immune response in the animal.

EXAMPLES

The present invention is illustrated by the following examples. It is to be understood that the particular examples, materials, amounts, and procedures are to be interpreted broadly in accordance with the scope and spirit of the invention as set forth herein.

Example I.

Purification and Characterization of a Cysteine Protease in the Murine Parasite *T. crassiceps*

A *T. crassiceps* cyst wall cysteine protease was purified to homogeneity (White et al., Mol. Biochem. Parasitol., 85:243-253 (1997)). The enzyme has an unusually high turnover ability for synthetic substrates and degrades human IgG rapidly *in vitro*. The molecule may play a key role in immune evasion by *Taenia*, a significant survival tactic employed by the parasite.

Because of its significant biological similarity to *T. solium*, rapid growth rates (> 100 cysts after 4 weeks of infection), and simple maintenance in BALB/c mice, *T. crassiceps* is commonly utilized as a model system for the study of the human parasite, *T. solium* (Hayunga et al., J. Parasitol. 75: 638-642 (1989); Ambrosio et al., Arch. Med. Res. 25:325-330 (1994); White et al., Infect. Agents Dis. 1:185-193 (1992)). *T. crassiceps*' natural hosts include rodents

(intermediate hosts) and canines (definitive hosts). We purified a predominant *T. crassiceps* cysteine protease to homogeneity, as confirmed by its N-terminus sequence, and its presentation as a single band on silver stained gels of sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). The enzyme was solubilized from whole parasites with acid extraction, whereas little cysteine protease activity was found in excretory fluid, a finding also noted by others (Hayunga et al., *J. Parasitol.* 75: 638-642 (1989); Ambrosio et al., *Arch. Med. Res.* 25:325-330 (1994)). Since acid extraction was able to dissociate most of the enzyme, in comparison to detergent solubilization, we hypothesized that the protease was located in the cyst wall, but was most likely not in the integral membrane. These observations were later supported by transmission electron microscopy, which localized cysteine protease activity to lysosomal vesicles and the cyst wall (Khalil et al., *J. Parasitol.* 84:513-515 (1998)). It is thus likely that this cysteine protease is located in the cyst wall of *Taenia*, rendering it accessible as a chemotherapeutic target.

Gel filtration (AcA 54 gel filtration column) and Fast Protein Liquid Chromatography (F.P.L.C.) anion exchange (Mono Q) chromatography were used to purify the cysteine protease 682-fold from the initial acid extract. The purified enzyme eluted at approximately 13% salt into 1 active sharp peak by anion exchange, and corresponded to 43 kDa based upon the expected molecular weight of the eluted fraction and the single band obtained by silver staining. (13% salt is significant since the similar *Taenia solium* cysteine protease also elutes at this percentage). Purity was further confirmed by an N-terminus sequence. Purity was further confirmed by an N-terminus sequence. Unambiguous results were obtained on the first 52 amino acids (Fig. 1(b)). An N-terminus sequence on material obtained from a second purification of *T. crassiceps* cysteine protease confirmed the sequence of the first 30 amino acids (Fig. 1a). Although the N-terminus sequence contains a cysteine at position 23, which is the catalytic cysteine in the papain superfamily, it does not contain the consensus pattern in enzymes of this superfamily (Hayunga et al., *J. Parasitol.* 75:

638-642 (1989)). However, the *T. crassiceps* cysteine protease most closely resembles enzymes in the papain family, in comparison to sequences of other cysteine protease families (Fig. 1b). Thus, this enzyme may represent a new class of cysteine proteases with greatest homology to the papain superfamily.

5 The purified cysteine protease was assayed for proteolytic activity using a synthetic peptide substrate linked to the C-terminus of 7-amino-4-trifluoromethyl coumarin (Z-Phe-Arg-AFC), substantially as described in White et al. (Mol. Biochem. Parasitol., 85:243-253 (1997)). Purified protease (0.1 ug) is added to a saturating amount of Z-Phe-Arg-AFC (1ug/500uL assay buffer) in
10 0.4M citrate supplemented with 10mM cysteine, at pH 4.9. The mixture is incubate for 18 hours at 37C. The purified cysteine protease was characterized by unequivocal thiol dependence (a biochemical characteristic of all cysteine proteases), inhibition by cysteine protease inhibitors, a pI of 5.27, and a substrate specificity profile demonstrating pronounced cleavage of Z-Phe-Arg-AFC (see
15 Example II). The purified protease was marked by an unusually high Kcat/Km (substrate turnover efficacy) on the synthetic peptide substrate, Z-Phe-Arg-AFC , equivalent to $2.2 \times 10^8 \text{ M}^{-1}\text{s}^{-1}$, which is an order of magnitude that approximates the enzymatic "diffusion control limit" which some investigators have likened to "catalytically perfect" enzymes (Voet, Biochemistry, 2nd ed, John Wiley and
20 Sons, eds. 14:371-400 (1995)). Very few enzymes have been reported with such a high catalytic Kcat/Km ratio. Most enzymes with these characteristics tend to be quick metabolic enzymes (e.g. hexokinase and triose phosphate isomerase), capable of instantly catalyzing reactions during every encounter with substrate. Thus, the impressive efficacy of the *Taenia* cysteine protease in turning over
25 substrate with similar magnitudes was suggestive of an enzyme with a biological significance to the parasite (such as metabolism).

The activity of the *T. crassiceps* protease was inhibited by the irreversible cysteine protease inhibitor, E64 (L-trans-epoxysuccinyl-leucylamide-(4-guanidino)-butane), indicating that the purified enzyme was a
30 cysteine protease. The enzyme's Km was substantially increased, indicating

substantial loss of affinity of the cysteine protease for Z-Phe-Arg-AFC substrate. This result is consistent with a covalent interaction between E64 and the purified enzyme, as is typically characteristic for the interaction of E64 and cysteine proteases. In order to obtain these kinetic parameters, assays were set up such that Km and Vm were calculated respectively from the x and y intercepts of a double reciprocal Lineweaver-Burke plot of substrate concentration (moles) versus protease velocity (mol h^{-1}). Kcat was calculated from the inverse of Vmax.

Example II.

Purification and Characterization of a Cysteine Protease in the Human Parasite, *T. solium*

A cysteine protease from the human parasite, *T. solium*, which uses the pig as an intermediate host, was purified and characterized. This protease is similar, although not identical, to the cysteine protease derived in Example I from *T. crassiceps*, whose natural hosts are rodents, and which grows very well in the BALB/c mouse. The data, however, are compelling that the *active sites* of the two proteases are highly homologous (which is the most important premise for a specific inhibitor). A protease inhibitor specifically directed against the *T. solium* cysteine protease is thus expected to specifically inhibit the cysteine protease in *T. crassiceps* thereby permitting testing for inhibitor efficacy in the mouse model.

T. solium cysts were obtained from wild cysticercotic pigs occupying the village of Guerrero in Mexico. As in *T. crassiceps*, most of the cysteine protease activity of the *T. solium* cysts could be released by acid extraction, without the need for detergent solubilization. When detergent solubilization was employed, the specific activity in the acid extracts was still approximately 5-fold higher. Thus, like the *T. crassiceps* cysteine protease, the *T. solium* cysteine protease is likely located in the cyst wall, but is not an integral membrane protein.

The purification protocol used in *T. crassiceps* was used to purify the *T.*

solium cysteine protease. After acid extraction, gel filtration chromatography was employed and the active fraction eluted at 50-55 kDa, thereby suggesting that the *T. solium* cysteine protease is slightly larger than the *T. crassiceps* cysteine protease (43 kDa). Optical density readings on FPLC anion exchange, showed that the profiles of the *T. solium* and *T. crassiceps* enzymes are practically superimposable (Fig. 2). Active fractions from the AcA 54 gel filtration column were fractionated further by FPLC anion exchange chromatography using a Mono Q column. The two enzymes were eluted with identical gradients (dashed line) of 10 mM Tris, pH 7.6 (solution A) and 10 mM Tris, pH 7.6 plus 1 M NaCl (solution B). One ml fractions were collected and assayed with Z-Phe-Arg-AFC substrate (dotted line). Absorbance was monitored at 280 nm (solid line). As can be observed, both enzymes elute between 12-14% salt, suggesting that these enzymes have a highly similar and conserved charge conformation. The *T. solium* cysteine protease electromigrated as a single band on silver stained SDS-PAGE gels.

These data suggest that there are substantial similarities between the *T. crassiceps* and *T. solium* cysteine proteases. Further evidence suggests that the active sites are homologous. First, the pH optimum of the *T. solium* cysteine protease is identical to the pH optimum of the *T. crassiceps* cysteine protease (pH 4.9). Moreover, the substrate cleavage patterns are similar. For both enzymes, Z-Phe-Arg-AFC (Z-FR-AFC) is the most significantly cleaved substrate in comparison to a panel of synthetic substrates (Fig. 3). Next to Z-FR-AFC, it can be seen that Lys-Ala-AFC (KA-AFC) and Arg-AFC (R-AFC) are the next best cleavable substrates by the *T. solium* protease. These two substrates are also the next best cleavable substrates for the *T. crassiceps* enzyme. The overall activity with respect to the other substrates (e.g., LG-AFC, ZRR-AFC) is not high enough in either enzyme. This data showing similar cleavage patterns between the two enzymes further confirms the homologous nature of the two proteases' active sites.

Like the *T. crassiceps* cysteine protease active site, the *T. solium* cysteine protease is unequivocally thiol dependent, and is completely inhibited by the absence of cysteine. As shown in Table 1, both cysteine proteases were inhibited by cysteine protease inhibitors but not significantly (>50%) by inhibitors of other catalytic classes. For example, inhibitors of the epoxy class of cysteine protease inhibitors (ie: E64 and EP459 (L-trans-epoxysuccinyl-leucylamido(4-methyl)butane)) completely abolish activity. HgCl₂ inhibits both enzymes in a concentration dependent manner, as is characteristic of cysteine proteases. Near total inhibition is noted with the chloromethane inhibitors tosyl lysyl chloromethyl ketone (TLCK) and tosylphenylanyl chloromethyl ketone (TPCK), which are known to inhibit serine proteases and some cysteine proteases. No significant inhibition is noted by inhibitors of other classes including PMSF, 1,10 phenanthroline, ethylenediaminetetracetic acid (EDTA) or pepstatin. In the table, one representative experiment is shown, but a minimum of three experiments were performed. Inhibitors were incubated with enzyme for 30 minutes before the addition of substrate. Fluorometric readings were assessed after overnight incubation.

Table 1. Sensitivity of (a) *T. crassiceps* (Example I) and (b) *T. solium* cysteine proteases to inhibition by a panel of inhibitors when Z-Phe-Arg-AFC is used as the substrate

a)

Inhibitor	Catalytic class	Concentration	% Inhibition \pm SD*
No cysteine			100
E-64	Cysteine	10^{-5} M	95 ± 3.0
		10^{-7} M	98 ± 1.5
		10^{-8} M	99 ± 1.0
Ep459	Cysteine	10^{-7} M	99 ± 1.1
Ep475	Cysteine	10^{-8} M	99 ± 0.9
Iodoacetate	Cysteine	10^{-3} M	99 ± 0.7
HgCl ₂	Cysteine	10^{-3} M	3 ± 2.1
		10^{-4} M	40 ± 29
		4×10^{-5} M	77 ± 24
PMSF	Serine	2×10^{-3} M	47 ± 32
APMSF	Serine	10^{-4} M	0 ± 0
TLCK	Serine/cysteine	10^{-4} M	96 ± 3.1
TPCK	Serine/cysteine	10^{-4} M	96 ± 3.5
1,10 Phenanthroline	Metallo	2×10^{-3} M	35 ± 38
EDTA	Metallo	2×10^{-3} M	27 ± 38
Peppstatin	Aspartic	10^{-6} M	34 ± 34

5

b)

Inhibitor*	Catalytic class	Concentration	% Inhibition \pm SD
No Cysteine			100
E-64	Cysteine	10^{-6} M	100 ± 2.7
		10^{-7} M	100 ± 0.2
		10^{-8} M	94 ± 0.9
Iodoacetate	Cysteine	10^{-7} M	100 ± 0.0
HgCl ₂	Cysteine	10^{-7} M	9 ± 3.1
		10^{-8} M	41 ± 10.0
		4×10^{-8} M	81 ± 1.4
PMSF	Serine	2×10^{-3} M	29 ± 10.0
TLCK	Serine/cysteine	10^{-4} M	93 ± 1.0
TPCK	Serine/cysteine	10^{-4} M	93 ± 1.0
1,10 Phenanthroline	Metallo	2×10^{-3} M	49 ± 10
EDTA	Metallo	1×10^{-3} M	13 ± 5.0
		2×10^{-3} M	8 ± 2.0
Peppstatin	Aspartic	1×10^{-5}	34 ± 4.0
		1×10^{-7}	39 ± 16
DCI			38 ± 11

10

Specific cysteine protease inhibitors (e.g., Suc-Leu-Tyr-AFC) abrogate the ability of the *T. solium* and *T. crassiceps* cysteine proteases to cleave human IgG on western blots (see Fig. 4b), when human IgG is used as a substrate. Thus, both enzyme active sites are marked by highly homologous inhibitor profiles.

In addition, the K_{cat}/K_m ratio for the *T. solium* cysteine protease active site, on Z-Phe-Arg-AFC is equal to $2.84 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ and is close to the K_{cat}/K_m ratio of the *T. crassiceps* cysteine protease ($2.2 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$). The *T. solium* cysteine protease is thus also within the logarithmic range of the diffusion control limit of 10^8 - 10^9 for "catalytically perfect" enzymes. Consequently, the *T. solium* cysteine protease also demonstrates a remarkable ability to turn over substrate, and we thus believe this to be evidence that such an efficacious enzyme must be significant to this human parasite.

Finally, like the *T. crassiceps* enzyme, the *T. solium* enzyme is also characterized by the capability to degrade the heavy and light chains of human IgG (Fig. 4), and the ability to do it quickly. Human IgG was incubated for 18 hours with a) *T. crassiceps* acid extract or b) *T. solium* purified cysteine protease (CP). Incubated tubes were subsequently fractionated by SDS-PAGE under reducing conditions, blotted onto PVDF membrane, and visualized using biotinylated anti-IgG (heavy chain/light chain), biotin-peroxidase conjugated-streptavidin complex and enhanced chemiluminescence. Top and bottom arrows indicate heavy and light chains, respectively, in panel A. Only the heavy chain is shown in Panel B. The absence of heavy and light chains suggests that the proteins are broken down into peptides. In another experiment, we noted that like *T. crassiceps*, the *T. solium* enzyme accomplishes complete IgG degradation by 30 minutes. This degradation was thiol dependent, and inhibited by the cysteine protease inhibitor, E64, but not by inhibitors of other catalytic classes. When the specific inhibitor, Z-LLY-FMK (Example IV)) was incubated with the *T. solium* cysteine protease in the presence of IgG, degradation of the antibody was inhibited. The *T. solium* cysteine protease also appears to be biased for human IgG degradation. For example, it did not degrade human albumin when this substrate was employed at the concentration that was used in the human IgG degradation studies.

- Collectively, our data suggest that the *T. solium* enzyme is biochemically and biologically nearly identical to the *T. crassiceps* enzyme, with regard to active site catalysis. Specific inhibitors that inhibit the *T. crassiceps* cysteine protease also inhibit the *T. solium* cysteine protease (Example IV). In summary,
- 5 evidence that supports the similarity of the enzyme active sites includes: highly similar pH optima, inhibitor profiles, substrate profiles, identical elution patterns off of anion exchange column, similar degradation of IgG, and inhibition by highly specific inhibitors. Consequently, the *T. crassiceps* cyst can confidently be used as a model system to study biological effects of protease inhibitors
- 10 against the *T. solium* cysteine protease.

Proposed interactions of Taenia cysteine proteases with the host's immune system

- The cyst wall cysteine proteases of *Taenia* may play a significant role in
- 15 *Taenia*'s evasion of host immunity. *Taenia* cysts survive for many years in asymptomatic intermediate hosts. Thus, *Taenia* cysts must somehow evade host immune responses. Moreover, they must find host protein sources to degrade in order to obtain essential amino acid nutrients. We postulate a mechanism of how this may occur, beginning with immune evasion. However, it is to be
- 20 understood that the present invention, insofar as it relates to methods and compounds for inhibiting the activity of a *Taenia* cysteine protease, is not intended to be limited by any particular mechanism or mode of action.

- It has previously been suggested that IgG traffics through the blood brain barrier or is produced in the brain. An Fc receptor has been identified on the
- 25 *Taenia* cyst wall, indicating that IgG uptake may be an active process (Kalinna et al., Parasitol. 106:289-296 (1993)). We postulate that IgG is routed to the cyst lysosomes in the cyst wall internuncial processes and tegumentary cytons. Pathways demonstrating protein shuttling to the lysosomes followed by protein degradation are known in *T. crassiceps* (Ambrosio et al., Arch. Med. Res. 25:325-330 (1994)). The cyst wall is where the cysteine protease is also
- 30

localized. After transport to the lysosomes, we hypothesize that the IgG would be degraded in the lysosomes. Hayunga et al. (Hayunga et al., J. Parasitol. 75: 638-642 (1989)) have observed degraded IgG units in *T. crassiceps* lysosomal vesicles. IgG is subsequently broken down by the specialized cysteine protease into peptides, or possibly even amino acids in the lysosomes. Thus, the IgG degrading cysteine protease may aid the cyst by protecting it from destructive host IgGs (complement-fixing IgG2a is associated to dying metacystodes).

Eventually, the benefits of accessibility to IgG -derived peptide or amino acids "lying around" in the lysosome may have become a significant metabolic advantage. Indeed, the cysteine protease may help the parasite actually *exploit* the host immune response, not just disable it. Due to the impressive efficiency of the cysteine protease's ability to degrade IgG, antibodies may have become a good source of nutrients for the parasite *in vivo*, as has been suggested (White et al., Mol. Biochem. Parasitol., 85:243-253 (1997)). The general importance of the cysteine protease in growing parasites is illustrated by our transmission electron microscope (TEM) studies (Khalil et al., J. Parasitol. 84:513-515 (1998)) which show greater cysteine protease activity in smaller, metabolically active and growing cysticerci than in larger, older ones. The correlation between increased cyst growth and increased cysteine protease activity in these smaller cysts is consistent with the suggestion that the enzyme may play a key role in breaking down proteins, like IgG, for parasite nutrition.

In the end, *Taenia* cysts may have become sensitized to a balance where a certain level of IgG in the host serum was actually beneficial metabolically, but too much could be destructive, immunologically. Perhaps there is a baseline physiological IgG level which is beneficial (IgG uptake in *T. crassiceps* cysts is shown to be saturable at physiological serum levels (Siebert et al., Exp. Parasitol. 48:64-174 (1979))). Thus, *Taenia* cysts may have a need for a certain level of IgGs for immune exploitation, but are harmed by concentration beyond this. Perhaps, this is why the cysteine protease, which is highly antigenic (Example III), is not located on the cyst wall surface, but is rather *within* the cyst

1 wall. Consequently, the cysts may have evolved molecular mechanisms to
control this balance. For example, a secretory *Taenia* glycoprotein (Villa et al.,
Parisitol. 112:561-570 (1996)) appears to modulate the shift from a host T
helper 1 (Th1) cellular mediated immune response to a humoral T helper 2
5 response (Th2), favoring increased antibody production which may be used by
the parasites for nutrition (this shift may also aid the parasites in avoiding a
destructive Th1 cellular response, for which they have no defenses).
Alternatively, diminished secretion of these molecules may slow down the shift
from Th1 to Th2 allowing the parasites another means to control their
10 immunological environment (and thus, physiological IgG levels).

We thus undertook to design or identify (Example IV) specific inhibitors
of *Taenia* cyst wall cysteine protease that potentially function using one or both
aspects of a dual mode of action. First, specific inhibition of the cysteine
protease may allow host immunoglobulins to recognize cysts thereby
15 "unmasking" them to the immune system. In Example VI, we show how cysts
from mice treated with specific inhibitors against the cysteine protease are
coated by immune cells whereas cysts from nontreated mice are "clean." A
second mode may be that inhibition of the cysteine protease may weaken the
cyst by depriving it of IgG derived amino acids, thus impairing it metabolically.
20 A "weakened" cyst would also more apt to be recognized by the immune system
and possibly destroyed or at least prevented from further growth and/or
proliferation. As shown in Example V our specific cysteine protease inhibitor
treatments protected mice from cysticercosis, suggesting that cyst growth (by
budding) was compromised.

25 Notably, the cysteine protease has been localized in these identical
regions in which the degraded IgG was reported. Transmission electron
microscopy (TEM) studies identified cysteine protease activity which cleaved
the cysteine protease substrate, Z-Phe-Arg-methoxynaphthylamide (MNA)
(Khalil et al., J. Parisitol. 84:513-515 (1998)) in acidic and lysosomal vacuoles
30 of the cyst wall in *T. crassiceps*. Free methoxynaphthylamide was coupled to

- osmium and p-rostanilin, and cleaved from Z-Phe-Arg at the substrate's carboxy side on the arginine residue. Compared to control without substrate, cysticeri incubated with substrate showed electron dense deposits in round vesicles, whereas most of these vesicles were found primarily within the cyst wall
- 5 tegumentary cytons and internuncial processes. This study thus localized the cysteine protease to the cyst wall. When the cysticeri were incubated with the cysteine protease inhibitor, E64, prior to substrate addition, the electron dense bodies were fewer and less dense than in cysticeri without the inhibitor. Thus, the cysteine protease activity here was partially inhibited by E64, further
- 10 supporting the presence of the cysteine protease in this region. It is hypothesized that inhibition was only partial with E64 because the concentration of this inhibitor within the acid vesicles may have been too low since the concentration of enzyme in the vesicles was higher than in our purification yields. Our important conclusions here are that 1) the cysteine protease appears
- 15 localized to the same cyst wall regions where degraded IgGs have been observed, and that 2) a cysteine protease inhibitor was able to reduce the cysteine protease's activity *in situ*. This latter observation suggests the promising possibility that specifically designed protease inhibitors may be able to permeate and consequently, inhibit cysteine protease activity in the *Taenia* cyst wall.
- 20 Moreover, the pH of the lysosomes (pH 4.9) is the identical pH optimum for activity of the cysteine protease on the synthetic substrate, Z-Phe-Arg-AFC, and the activity of the cysteine protease is totally dependent upon exogenous reducing groups, which are also present in the lysosomes. Although Z-Phe-Arg-AFC is cleaved by cysteine proteases as well as by serine proteases
- 25 like kallikreins, serine proteases do not optimally function at acidic pH and they are thiol dependent. Also, prior studies failed to identify serine protease activity in *Taenia* cysticeri walls with similar peptides. Thus, the proteolytic activity detected here with Z-Phe-Arg-AFC at acidic pH in the reducing environment of the lysosomes is most likely due to the cysteine protease that we purified from

the *Taenia* cyst wall. We also observed that the cysteine protease which we purified is the only one in the cyst wall.

The *Taenia* cysteine proteases cleave human IgG optimally at lysosomal pH and only in the presence of reducing groups, *in vitro*. It is thus notable that this enzyme was solely responsible for complete human IgG digestion on western blots, in comparison to other proteases present in the original acid extract of the purification protocol. The observation that both the heavy chain and light chains could be completely cleaved by the cysteine protease, suggests that the chains were reduced to peptides or amino acids, which are not visible on SDS-PAGE gels. Dose response experiments showed that the purified cysteine protease completely degraded both heavy and light chains of human IgG at a molar ratio up to 1:30 (enzyme: IgG), whereas partial degradation of IgG occurred at molar ratios up to 1:3000. Complete degradation was accomplished within only 45 minutes and was completely inhibited by the cysteine protease inhibitor, E64. As mentioned, E64 suppresses cysteine protease activity in the *Taenia* cyst wall *in situ*, suggesting that specifically designed cysteine protease inhibitors should also suppress IgG degradation by the cysteine protease. (Indeed, specific inhibition of the homologous *T. solium* cysteine protease is noted for Leu-Tyr-AFC-Fig 4b). Moreover, the substrate specificity profile of the purified enzyme most closely resembles cathepsin L-like cysteine proteases, which are mammalian lysosomal enzymes, and have also been linked to IgG degradation. These data are therefore compelling that the *T. crassiceps* cysteine protease efficiently degrades host IgG substrates.

Our data also suggest that the *Taenia* cysteine protease may also be preferentially selective for IgG degradation. For example, we have recently isolated a secretory cysteine protease from *T. crassiceps*, with a pH optimum of 6.3, which appears to cause only minor cleavage of the human IgG heavy chain but not the light chain. By comparison, the cyst wall cysteine protease appears to degrade both chains. Based upon these data, we believe that the *Taenia* cyst

wall cysteine protease significantly contributes to lysosomal degradation of host IgG in *Taenia*, and may also be specialized for this process.

In summary, the *Taenia* cyst wall cysteine protease's unusually high kinetic turnover ratio, its potentially key role in parasite nutrition and host immune evasion, its accessible location within the cyst wall (but not too deep within the cyst), and the fact that its activity can be reduced by cysteine protease inhibitors *in situ*, render this enzyme a promising target for antiparasitic chemotherapy.

Example III.

Vaccination of BALB/c Mice with the *T. solium* Cysteine Protease

BALB/c mice were vaccinated with the purified (eluted off the MonoQ anion exchange column) and partially purified (eluted off ACA 54 gel filtration column) *T. solium* cysteine protease (Example II) in Freund's Complete Adjuvant. Mice were subsequently boosted with the same preparations two weeks later. Five and a half months following the booster, the mice were challenged with *T. crassiceps* cysts followed by euthanization after 6 weeks of active infection. As can be seen in Table 2 below, mice vaccinated with the purified cysteine protease were protected 72% from cysticercosis, whereas mice vaccinated with the partially purified preparation, were protected 50%. The *T. solium* cysteine protease thus appears to be able to induce a cross-protective immune response in BALB/c mice, and this response correlates to the purity of the cysteine protease. This observation supports the characterization of these enzymes as antigenically similar.

Table 2. Protection of BALB/c mice from *T. crassiceps* cystercercosis infection, after immunization with the purified (ACA54 followed by Mono Q) and partially purified (ACA54 only) *T. solium* cysteine proteinase

5

	Cyst #	AVG	STD	% Protecti on	Cyst Volum e	AVG	STD
<i>MonoQ</i>	43	38.6	18.6	71.8	1.6	1.3	.6
<i>(pure)</i>	59				1.8		
	14				.5		
<i>ACA54</i>	62	69	5.3	49.8	2	2.1	.12
<i>(partially</i>	75				2.3		
<i>pure)</i>	70				2.1		
<i>Positive</i>	158	137.	18.6		7	7.2	.21
	3						
<i>(no</i>	113				7.5		
<i>vaccine)</i>	141				7.1		

We also obtained splenocytes from *T. crassiceps* infected mice and stimulated them with the purified *T. solium* cysteine protease or with a *T. solium* extract. Splenocytes were removed from mice carrying cysticercosis infection for 6 weeks as well as from normal, non-infected mice of the same age. The normal cells were exposed to Concanavalin-A (Con-A). The cells from the infected mice were exposed to the purified *T. solium* cysteine protease and the starting *T. solium* extract. Cellular proliferation was measured by ³H-thymidine incorporation. The purified protease elicited a proliferative response in the immune cells that was significantly greater than that elicited by the extract, equaling stimulation by the mitogen Concanavalin-A (Fig. 5).

Example IV.

Identification of Bound, Potent, and Selective Inhibitors Against the *T. crassiceps* and *T. solium* Cysteine Proteases *in Vitro*

It was shown in Example II that the cyst wall proteases from *T. solium* and *T. crassiceps* are biochemically similar. Thus, a protease inhibitor directed against the *T. solium* cyst wall cysteine protease would be expected to inhibit the cyst wall cysteine proteinase of *T. crassiceps*.

Our results with a Langmuir binding isotherm revealed that the *T. solium* cysteine proteinase has only one substrate binding site, suggesting that designing a specific inhibitor was feasible. Our goal in this study was to demonstrate efficacious, potent and selective inhibition of the enzyme *in vitro*, preferably characterized by binding of the inhibitor at the active site. Inhibitors that bind at the active site of an enzyme can also be used to assess the similarities of the enzymes' active sites (Fig. 6). Three inhibitors of *Taenia* cyst wall cysteine protease were identified, and the protective potential of two of these inhibitors *in vivo* in the treatment of murine cysticercosis was also evaluated.

Two protease inhibitors inhibited the purified *T. solium* cysteine proteases in an assay using Z-Phe-Arg-AFC as a substrate at low efficacious concentrations: Z-Leucine-Leucine-Leucine-Fluoromethylketone (Z-LLL-FMK) and Z-Leucine-Leucine-Tyrosine-Fluoromethylketone (Z-LLY-FMK). At 5 nanomolar levels, these protease inhibitors suppressed the *T. solium* cysteine protease's activity by 100% and 97%, respectively. Our kinetic data show that these inhibitors were bound to the cysteine protease's active site and were also potent for its inactivation. A third inhibitor, Ph2-LF-KP was also noted to inhibit the target cysteine protease rather efficaciously, by 80% when used at the 10 same concentration as Z-LLL-FMK and Z-LLY-FMK. The inhibitors (Z-LLL-FMK and Z-LLY-FMK) similarly inhibited the *T. crassiceps* cysteine protease.

Z-LLY-FMK and Z-LLL-FMK also inhibited degradation of human IgG by the *T. solium* cysteine protease. Thus, we hypothesized that they may also have the potential to block human IgG degradation by *Taenia* cysts *in vivo*. This 15 consideration, the inhibitors' impressive efficacy in inhibiting the target proteases at low concentrations, and their nontoxicity to immune cells *in vitro* (Example VII) suggest that these inhibitors would serve as good candidates for chemotherapy in the mouse model of cysticercosis.

20 Example V.

Protection of BALB/c Mice from Cysticercosis Infection using the Specific Inhibitors Z-Leucine-Leucine-Tyrosine Fluoromethylketone (Z-LLY-FMK)

and

Z-Leucine-Leucine-Leucine-Fluoromethylketone (Z-LLL-FMK)

25 Based upon the premise that the cyst wall cysteine protease may serve a critical function to the life cycle of the *Taenia* cyst, we tested our most efficacious inhibitors, Z-LLL-FMK, Z-LLY-FMK and Ph2-LF-KP for their abilities to protect mice from cysticercosis infection. Z-LLY-FMK and Z-LLL-FMK were individually tested as prophylactics in a preliminary trial with 30 BALB/c mice. Injections were carried out in 150 μ l of dimethyl sulfoxide. Mice

were pre-injected intraperitoneally for two days with the inhibitor ($\sim 1.4 \times 10^{-2}$ M, dissolved in an injection vol. of 150 μ l 0.15M PBS) followed by infection with 10 *T. crassiceps* cysts/mouse in 200 μ l 0.15 M PBS. Mice were subsequently dosed daily with the same concentration for four weeks. After one month of infection, both groups were euthanized and cysts removed by washing with sterile 0.15 M PBS. A minimum of two attached cysts were scored as multilobed (ML). Upon visual inspection, no mice showed effects of inhibitor toxicity.

Mice in groups treated with Z-LLL-FMK were protected 100% from cysticercosis infection (Table 3). Mice in groups treated with Z-LLY-FMK were protected 85%-97% from cysticercosis infection, in comparison to untreated controls (Tables 3 and 4). Mice in groups treated with Ph2-LF-KP were protected 40 %. A subsequent study repeated the identical prophylactic experiment with Z-LLY-FMK and demonstrated similar protection data (75%-90%).

Table 3. Correlation between *in vitro* inhibition of *T. solium* cysteine protease and *in vivo* protection

<i>Inhibitor</i>	% <i>Inhibition</i> of <i>T. solium</i> CP <i>in vitro</i>	% <i>Prophylactic</i> protection of BALB/c mice	% <i>Therapeutic</i> protection of BALB/c mice	<i>Immune cells</i> observed by SEM on cyst surfaces
Z-LLL-FMK	100%	100%	60%	N/a
Z-LLY-FMK	97%	85-97%	n/a	Intense
Ph2-LF-KP	80%	40%	n/a	None observed

"n/a" refers to unavailable data.

Table 4. The effect of Z-Leu-Leu-Tyr-Fluoromethylketone treatment of BALB/c mice challenged with *Taenia crassiceps* cysts

Untreated controls	Cyst #	Avg	SD	% Prot.	# ML	% ML	Avg
Mouse 1	240	197.7	34.3		30	12.5 %	14.4%
Mouse 2	156				16	10.3%	
Mouse 3	197				40	20.3%	
Treated controls							
Mouse 4	7	21.3	10.2	96.5%	4	57%	51%
Mouse 5	30			84.8%	13	43%	
Mouse 6	27			86.4%	14	52%	

- 5 Interestingly, a high percentage of cysts which did survive the treatment demonstrated abnormal morphology. Many of these cysts exhibited apoloar multilocularity (multi-lobed appearances), an abnormal budding pattern. Under histological exam, multilobed cysts showed enlarged walls compared to normal lobed cysts.
- 10 A therapeutic study with Z-LLL-FMK was also conducted. During treatment phase, each treated mouse received 1.48×10^{-2} ug/ul of Leu-Leu-Tyr-FMK daily for three weeks and every other day during the last week. The results revealed 60% protection when inhibitor treatment began two weeks after mice had been infected with *T. crassiceps* cysts (Table 3). Mice in all groups survived
- 15 without noticeable side effects (e.g., coat condition, tail dragging or paresis, etc).
- Finally, it is notable that a dose response of *in vitro* inhibition of the *T. solium* cysteine protease correlates with *in vivo* protection in BALB/c mice. Moreover, cysts removed from mice treated with the less effective inhibitor, Ph2-LF-KP, had neither adherent immune cells nor damaged tegument.

Example VI. Morphological Evaluation of Cysts in Mice Treated with Inhibitor of *Taenia* Cysteine Protease Activity

Scanning electron microscopy (SEM) examination of the few surviving cysts which were removed from mice treated with Z-LLY-FMK in Example V revealed the presence of adherent immune cells, which were concomitantly present where the cyst wall appears to be undergoing destruction (Fig. 7). We identified macrophages, neutrophils, fibroblasts, and collagenous deposits on the cyst surfaces from treated mice, in comparison to their absence on cysts from untreated mice. General surface atrophy was also apparent. Microtriches (microvilli) were either sloughed off in entire regions or significantly shortened in general, compared to cysts which were removed from untreated mice. Breaks in the integrity of the cyst surface were also noted for the cysts removed from treated mice. Higher magnification revealed tegumental erosion near immunological cells. Cysts removed from untreated mice were characterized by no immune cells, intact and longer microtriches, and homogeneity of the cyst surface. These results are consistent with other studies which have shown that viable cysticerci in pig muscle and those removed from humans at autopsy show little surrounding host inflammation.

Example VII.

Z-LLY-FMK and Z-LLL-FMK are Nontoxic to Splenocytes from BALB/c Mouse Cells *In Vitro*

Mouse splenocytes were removed and treated with either inhibitor alone or inhibitor and concanavalin A (Con-A). Splenocytes from normal mice were removed and treated with either inhibitor alone, or inhibitor and concanavalin A. Proliferation was measured by ³H-thymidine incorporation into the cells. A complete dose response curve for the inhibitor was conducted, with the highest dose representing in Fig. 8. The inhibitors did not induce a proliferative response (as measured by ³H-thymidine incorporation). Furthermore, the inhibitors did not affect the responsiveness of the splenocytes to Con-A.

Moreover, this dosage is three times the concentration which was utilized to inject mice on a daily basis in the prophylactic experiments (Example V).

Furthermore, trypan blue staining demonstrated that the mouse cells were all viable, even if the inhibitor was present. These studies further support our observation that no host side effects were observed in mice when they were treated with the specific inhibitors, *in vivo*. Z-LLL-FMK and Z-LLY-FMK thus appear to be nontoxic to normal BALB/c mouse splenocytes *in vitro*.

Example VIII.

Delivery of *Taenia* Cysteine Protease Inhibitor to Humans

Inhibitors for medical or veterinary use are designed to inhibit the activity of *Taenia* cysteine protease in humans, in pigs or in both. In pigs, *T. solium* cysts are located predominantly in muscles. However, the parasite is very neurotropic in humans, hence the disease often manifests itself as neurocysticercosis. In neurocysticercosis the *T. solium* cysts are located in the brain, thus it is desirable to design inhibitors that are able to cross the blood-brain barrier. The murine *T. crassiceps* model is well-suited for determining the *in vivo* susceptibility of tapeworm larvae (cysts) to drugs, but potential drugs for human use also need to be evaluated for their ability to cross the blood-brain barrier. Drugs like albendazole and praziquantel that affect brain cysts are surmised to cross the barrier.

It has been experimentally shown that the oil-water partition coefficient of nonelectrolytes is correlated with their membrane permeability and brain uptake in animal models (Levin, J. Med. Chem., 23:682-684(1980); Cornford, Exper. Parasitol., 70:25-34 (1990)). Therefore, partition coefficients for the inhibitor compounds will be determined, and if their values predict barrier impermeability, strategies are pursued to improve the likelihood of permeation. A variety of strategies for by-passing this barrier are known in the art, involving chemical modification of drugs, physical alteration of the blood/brain barrier and circumvention of the blood-brain barrier (reviewed in Abbott et al., Mol. Med.

Today, March 1996, 106-113). For example, Alkermes, Inc. of Cambridge, MA, a company that specializes in drug-delivery systems, has a promising blood-brain barrier permeabilizing drug, RMP-7 (Cereport®), that is now in Phase II of clinical trials. As another example, the candidate drugs can be are chemically
5 modified to improve their partition coefficients. The FMK substituent, for example, is thought to be important in membrane permeability of the drug and may assist in transporting the drug across the blood-brain barrier.

The complete disclosures of all patents, patent applications including
10 provisional patent applications, and publications, and electronically available material (e.g., GenBank amino acid and nucleotide sequence submissions) cited herein are incorporated by reference. The foregoing detailed description and examples have been provided for clarity of understanding only. No unnecessary
15 limitations are to be understood therefrom. The invention is not limited to the exact details shown and described; many variations will be apparent to one skilled in the art and are intended to be included within the invention defined by the claims.

WHAT IS CLAIMED IS:

1. An isolated cyst wall cysteine proteinase derived from *Taenia solium*.
2. An isolated polynucleotide having a nucleic acid sequence that encodes the proteinase of claim 1.
3. An isolated polynucleotide that is the complement of the nucleic acid molecule of claim 2.
4. A cyst wall cysteine proteinase or a polynucleotide having a nucleic acid sequence encoding a cyst wall cysteine proteinase, for use as a vaccine.
5. Use of a cyst wall cysteine proteinase or a polynucleotide having a nucleic acid sequence encoding a cyst wall cysteine proteinase for the preparation of a vaccine.
6. A vaccine comprising at least one component selected from the group consisting of (a) a polypeptide comprising a cyst wall cysteine proteinase or an immunogenic polypeptide subunit thereof and (b) a polynucleotide comprising a nucleotide sequence encoding a polypeptide comprising a cyst wall cysteine proteinase or an immunogenic polypeptide subunit thereof.
7. The vaccine of claim 6 wherein the cyst wall cysteine proteinase is derived from *T. solium* or *T. crassiceps*.

8. A method for treating an animal harboring a *Taenia* infection comprising administering to the infected animal the vaccine of claims 6 or 7, wherein administration of the vaccine is effective to eliminate the parasite from the animal or to prevent or delay the appearance of cysticercosis or neurocysticercosis the animal.

9. A method for protecting an animal against a *Taenia* infection comprising administering to an uninfected animal the vaccine of claims 6 or 7, wherein administration of the composition is effective to prevent subsequent infection of the animal by the parasite or to prevent the development of cysticercosis or neurocysticercosis in the animal after subsequent infection by the parasite.

10. The method of claims 8 or 9 wherein the animal is a pig or a human and the *Taenia* infection is a *T. solium* infection.

11. A pharmaceutical composition comprising:
an inhibitor molecule that inhibits the activity of a cyst wall
cysteine protease, the inhibitor molecule comprising a peptide or
peptidomimetic compound; and
a pharmaceutically acceptable carrier.

12. The pharmaceutical composition of claim 11 wherein the peptide or peptidomimetic compound comprises (Xaa)_n-Yaa-Zaa-R; wherein Xaa and Zaa are each independently any amino acid; Yaa is a hydrophobic amino acid; R comprises a nucleophilic moiety; and n = 0-5.

13. The pharmaceutical composition of claim 12 wherein Yaa is leucine.

14. The pharmaceutical composition of claims 12 or 13 wherein R comprises a chemical moiety selected from the group consisting of a carboxylic acid derivative, an amide derivative, a benzene derivative, a phenyl derivative, a chloromethylketone or derivative thereof, a fluoromethylketone or derivative thereof, an alphaketo acid or derivative thereof, a ketoamide or derivative thereof, a ketoester or derivative thereof, a vinylsulfone or derivative thereof, and a pyridyl or derivative thereof.

15. The pharmaceutical composition of claims 12-14 wherein R comprises a fluoromethylketone (FMK).

16. The pharmaceutical composition of claim 15 wherein the inhibitor molecule is Z-Leu-Leu-Leu-FMK or Leu-Leu-Tyr-FMK.

17. The pharmaceutical composition of claim 11-16 wherein the inhibitor molecule inhibits cyst wall cysteine proteinase derived from *T. solium* or *T. crassiceps*.

18. Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK for use as an active pharmaceutical substance.

19. Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK according to claim 18 for use in treating human neurocysticercosis or porcine cysticercosis.

20. The use of Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK for the preparation of a pharmaceutical composition.

21. The use of an inhibitor molecule that inhibits the activity of a *Taenia* cysteine protease, the inhibitor molecule comprising a peptide or peptidomimetic compound, for the treatment of human neurocysticercosis or porcine cysticercosis.

22. The use of an inhibitor molecule according to claim 21 wherein the peptide or peptidomimetic compound comprises $(Xaa)_n$ -Yaa-Zaa-R; wherein Xaa and Zaa are each independently any amino acid; Yaa is a hydrophobic amino acid; R comprises a nucleophilic moiety; and $n = 0-5$.

23. The use of an inhibitor molecule according to claim 22 wherein Yaa is leucine.

24. The use of an inhibitor molecule according to claims 22 or 23 wherein R comprises a chemical moiety selected from the group consisting of a carboxylic acid derivative, an amide derivative, a benzene derivative, a phenyl derivative, a chloromethylketone or derivative thereof, a fluoromethylketone or derivative thereof, an α -keto acid or derivative thereof, a ketoamide or derivative thereof, a ketoester or derivative thereof, a vinylsulfone or derivative thereof, and a pyridyl or derivative thereof.

25. The use of an inhibitor molecule according to claims 22-24 wherein R comprises a fluoromethylketone (FMK).

26. A method for inhibiting *Taenia* cyst wall cysteine protease activity comprising contacting a *Taenia* cyst wall cysteine protease with an inhibitor molecule comprising $(Xaa)_n$ -Yaa-Zaa-R; wherein Xaa and Zaa are each any amino acid; Yaa is a hydrophobic amino acid; R comprises a nucleophilic moiety; and n is 0 to about 5.

27. The method of claim 26 wherein Yaa is leucine.
28. The method of claims 26 or 27 wherein R comprises a chemical moiety selected from the group consisting of a carboxylic acid derivative, an amide derivative, a benzene derivative, a phenyl derivative, a chloromethylketone or derivative thereof, a fluoromethylketone or derivative thereof, an alphaketo acid or derivative thereof, a ketoamide or derivative thereof, a ketoester or derivative thereof, a vinylsulfone or derivative thereof, and a pyridyl or derivative thereof.
29. The method of claim 26-28 wherein R comprises a fluoromethylketone (FMK).
30. The method of claim 29 wherein the inhibitor molecule is Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK.
31. The method of claim 26-30 performed in cell free environment.
32. The method of claim 26-30 performed in cell culture, in an organ, or in a tissue.
33. The method of claim 26-30 performed in a whole animal.
34. The method of claim 26-30 wherein the *Taenia* cysteine proteinase is derived from *T. solium*.

35. A method for identifying an inhibitor of *Taenia* cysteine proteinase activity comprising:

combining a candidate inhibitor and a *Taenia* cysteine proteinase to form a mixture;

adding the proteinase substrate Z-Phe-Arg-7-amino-4-trifluoromethylcoumarin to the mixture; and

determining the extent to which the proteinase substrate is cleaved; wherein a reduction in the extent of substrate cleavage compared to the extent of substrate cleavage in the absence of the candidate inhibitor is indicative of inhibition of *Taenia* cysteine proteinase activity.

36. The method of claim 35 comprising combining a candidate inhibitor and a cysteine protease derived from *T. solium* or *T. crassiceps* to form the mixture.

37. A computer-assisted method for identifying an inhibitor of *Taenia* cysteine proteinase activity comprising:

supplying a computer model of the structure of an inhibitor of *Taenia* cysteine proteinase activity;

either (i) computationally or visually screening a structural library for candidate compounds having a structure similar to that of the inhibitor or (ii) computationally or visually designing a candidate compound having a structure similar to that of the inhibitor; and

assaying the candidate compound for the ability to inhibit *Taenia* cysteine proteinase activity.

38. A computer-assisted method for identifying an inhibitor of *Taenia* cysteine proteinase activity comprising:

solving the X-ray crystal structure of a *Taenia* cysteine proteinase to yield a computer model of the *Taenia* cysteine proteinase;

supplying a computer model of the structure of an inhibitor of *Taenia* cysteine proteinase activity;

computationally or visually docking the inhibitor structure to the proteinase crystal structure at the binding site of the inhibitor;

computationally or visually identifying intermolecular interactions between the inhibitor and the proteinase;

either (i) computationally or visually screening a structural library for candidate compounds having a structure similar to that of the inhibitor or (ii) computationally or visually designing a candidate compound having a structure similar to that of the inhibitor;

computationally or visually evaluating the intermolecular interactions between the candidate compound and the proteinase; and

assaying the candidate compound for the ability to inhibit *Taenia* cysteine proteinase activity.

39. The method of claims 37 or 38 wherein the inhibitor is Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK.

40. A method for treating an animal harboring a *Taenia* infection comprising administering to the infected animal a pharmaceutical composition comprising an inhibitor of a *Taenia* cysteine proteinase, wherein administration of the composition is effective to eliminate the parasite from the animal or to prevent or delay the appearance of cysticercosis or neurocysticercosis in the animal.

41. A method for protecting an animal against a *Taenia* infection comprising administering to an uninfected animal a pharmaceutical composition comprising an inhibitor of a *Taenia* cysteine proteinase, wherein administration of the composition is effective to prevent subsequent infection of the animal by the parasite or to prevent the development of cysticercosis or neurocysticercosis the animal after subsequent infection by the parasite.

42. The method of claims 40 or 41 wherein inhibitor is Z-Leu-Leu-Leu-FMK or Z-Leu-Leu-Tyr-FMK.

43. The method of claims 40, 41 or 42 wherein the animal is a pig or a human and the *Taenia* infection is a *T. solium* infection.

10/018936

1/9

A.

Position 1 6 11 16 17 18 19 20 21 22 23 24

Amino Acid GLKET VEKLF DEFDK DKSGK ISCAE LKAAL

(SEQ ID NO:1)


B.

	15	16	17	18	19	20	21	22	23	24
(SEQ ID NO:2) <i>I. crassiceps</i> cysteine proteinase	K	D	K	S	G	K	I	S	C	A
(SEQ ID NO:3) <i>Fasciola</i> cathepsin L	K	D	Q	G	Q	C	G	S	C	W
(SEQ ID NO:4) <i>Schistosoma mansoni</i> cathepsin B	R	D	Q	S	R	C	G	S	C	W
(SEQ ID NO:5) Papain	K	N	Q	G	S	C	G	S	C	W
(SEQ ID NO:6) <i>Paragonimus</i> cysteine proteinase	E	N	Q	G	E	C	G	S	C	W
(SEQ ID NO:7) <i>Schistosoma mansoni</i> calpain	-	C	Q	G	E	L	G	D	C	W
(SEQ ID NO:8) <i>Schistosoma mansoni</i> hemoglobinase	N	Y	R	H	Q	A	D	V	C	H
(SEQ ID NO:9) IL-1 β converting enzyme	K	P	K	V	I	I	Q	A	C	R
(SEQ ID NO:10) <i>Streptococcus</i> cysteine proteinase	V	G	Q	A	A	T	G	H	C	V

Fig. 1


10/018936

2/9

*Fig. 2a*

10/018936

3/9

*Fig. 2b*

4/9

10/018936




Fig. 3a




Fig. 3b

10/018936

5/9




Fig. 4

10/018936

6/9





Fig. 5

10/018936

7/9

(Ph)₂CHCO-Leu-Phe-CO-NHCH₂-2-Pyridyl - (Ph2-LF-KP)

Z-Leu-Phe-CONH-Ethyl - (Z-LF-KE)

Z-Leu-Phe-CONH-4-Morphophinyl - (Z-LF-KM)

Z-Leu-Phe-CH₂-CH(OH)-C₆H₄-(4-Oph)- (Z-LF-BPh) [B=benzene]

Boc-Leu-Phe-VinylSulfone-Phenyl - (B-LF-VS)

Z-Leu-Leu-Leu-VinylSulfone-Phenyl - (Z-LLL-VS)

Z-Leu-Leu-Tyr-Fluoromethylketone (Z-LLY-FMK)

Fig. 6

10/018936

8/9




Fig. 7

10/018936

9/9




Fig. 8

DECLARATION AND POWER OF ATTORNEY

We, Salman Baig, Raymond T. Damian, and Arthur Clinton White, declare that: (1) our respective citizenships and residence/mailling addresses are indicated below; (2) we have reviewed and understand the contents of the specification identified below, including the claims, as amended by any amendment specifically referred to herein. (3) we believe that we are the original, first, and joint inventors of the subject matter in

**CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF
NEUROCYSTICERCOSIS**

International Patent Application Serial No. PCT/US00/10672, filed 20 April 2000

described and claimed therein and for which a patent is sought; and (4) we hereby acknowledge our duty to disclose to the United States Patent and Trademark Office all information known to us to be material to the patentability as defined in Title 37, Code of Federal Regulations, §1.56.

We hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate listed below, or §365(a) of any PCT international application which designates at least one country other than the United States of America listed below, and have also identified below any foreign application for patent or inventor's certificate, or any PCT international application having a filing date before that of the application on the basis of which priority is claimed:

a. ☒ no such applications have been filed.

b. ☐ such applications have been filed as follows:

FOREIGN APPLICATION(S), IF ANY, CLAIMING PRIORITY UNDER 35 USC §119(a)-(d), §365(a), and/or §365(b)			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)

ALL FOREIGN APPLICATIONS, IF ANY, FILED BEFORE THE PRIORITY APPLICATION(S)			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)

DECLARATION AND POWER OF ATTORNEY

We, Salman Baig, Raymond T. Damjan, and Arthur Clinton White, declare that: (1) our respective citizenships and residence/mailling addresses are indicated below; (2) we have reviewed and understand the contents of the specification identified below, including the claims, as amended by any amendment specifically referred to herein, (3) we believe that we are the original, first, and joint inventors of the subject matter in

CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

International Patent Application Serial No. PCT/US00/10672, filed 20 April 2000

described and claimed therein and for which a patent is sought; and (4) we hereby acknowledge our duty to disclose to the United States Patent and Trademark Office all information known to us to be material to the patentability as defined in Title 37, Code of Federal Regulations, § 1.56.

We hereby claim foreign priority benefits under Title 35, United States Code, § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate listed below, or § 365(a) of any PCT international application which designates at least one country other than the United States of America listed below, and have also identified below any foreign application for patent or inventor's certificate, or any PCT international application having a filing date before that of the application on the basis of which priority is claimed:

- a. ☒ no such applications have been filed.
 b. ☐ such applications have been filed as follows:

FOREIGN APPLICATION(S), IF ANY, CLAIMING PRIORITY UNDER 35 USC § 119(a)-(d), § 365(a), and/or § 365(b)			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)

ALL FOREIGN APPLICATIONS, IF ANY, FILED BEFORE THE PRIORITY APPLICATION(S)			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)

DECLARATION AND POWER OF ATTORNEY

We, Salman Baig, Raymond T. Damian, and Arthur Clinton White, declare that: (1) our respective citizenships and residence/mailing addresses are indicated below; (2) we have reviewed and understand the contents of the specification identified below, including the claims, as amended by any amendment specifically referred to herein, (3) we believe that we are the original, first, and joint inventors of the subject matter in

CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

International Patent Application Serial No. PCT/US00/10672, filed 20 April 2000

described and claimed therein and for which a patent is sought; and (4) we hereby acknowledge our duty to disclose to the United States Patent and Trademark Office all information known to us to be material to the patentability as defined in Title 37, Code of Federal Regulations, §1.56.*

We hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate listed below, or §365(a) of any PCT international application which designates at least one country other than the United States of America listed below, and have also identified below any foreign application for patent or inventor's certificate, or any PCT international application having a filing date before that of the application on the basis of which priority is claimed:

a. ☒ no such applications have been filed.

b. ☐ such applications have been filed as follows:

FOREIGN APPLICATION(S), IF ANY, CLAIMING PRIORITY UNDER 35 USC §119(a)-(d), §365(a), and/or §365(b)			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)

ALL FOREIGN APPLICATIONS, IF ANY, FILED BEFORE THE PRIORITY APPLICATION(S)			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)

* Title 37, Code of Federal Regulations, §1.56 is reproduced on the attached page.

Applicant(s): Baig et al.

Serial No.: PCT/US00/10672

Int'l Filing Date: April 20, 2000

Title: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

§ 1.56 Duty to disclose information material to patentability.

(a) A patent by its very nature is affected with a public interest. The public interest is best served, and the most effective patent examination occurs when, at the time an application is being examined, the Office is aware of and evaluates the teachings of all information material to patentability. Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is cancelled or withdrawn from consideration, or the application becomes abandoned. Information material to the patentability of a claim that is cancelled or withdrawn from consideration need not be submitted if the information is not material to the patentability of any claim remaining under consideration in the application. There is no duty to submit information which is not material to the patentability of any existing claim. The duty to disclose all information known to be material to patentability is deemed to be satisfied if all information known to be material to patentability of any claim issued in a patent was cited by the Office or submitted to the Office in the manner prescribed by §§1.97(b)-(d) and 1.98. However, no patent will be granted on an application in connection with which fraud on the Office was practiced or attempted or the duty of disclosure was violated through bad faith or intentional misconduct. The Office encourages applicants to carefully examine:

- (1) Prior art cited in search reports of a foreign patent office in a counterpart application, and
- (2) The closest information over which individuals associated with the filing or prosecution of a patent application believe any pending claim patentably defines, to make sure that any material information contained therein is disclosed to the Office.

(b) Under this section, information is material to patentability when it is not cumulative to information already of record or being made of record in the application, and

- (1) It establishes, by itself or in combination with other information, a *prima facie* case of unpatentability of a claim; or
- (2) It refutes, or is inconsistent with, a position the applicant takes in:
 - (i) Opposing an argument of unpatentability relied on by the Office, or
 - (ii) Asserting an argument of patentability.

A *prima facie* case of unpatentability is established when the information compels a conclusion that a claim is unpatentable under the preponderance of evidence, burden-of-proof standard, giving each term in the claim its broadest reasonable construction consistent with the specification, and before any consideration is given to evidence which may be submitted in an attempt to establish a contrary conclusion of patentability.

(c) Individuals associated with the filing or prosecution of a patent application within the meaning of this section are:

- (1) Each inventor named in the application;
- (2) Each attorney or agent who prepares or prosecutes the application; and
- (3) Every other person who is substantively involved in the preparation or prosecution of the application and who is associated with the inventor, with the assignee or with anyone to whom there is an obligation to assign the application.

(d) Individuals other than the attorney, agent or inventor may comply with this section by disclosing information to the attorney, agent, or inventor.

(e) In any continuation-in-part application, the duty under this section includes the duty to disclose to the

Office all information known to the person to be material to patentability, as defined in paragraph (b) of this section, which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

Declaration and Power of Attorney

Applicant(s): Baig et al.

Serial No.: PCT/US00/10672

Int'l Filing Date: April 20, 2000

Title: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

We hereby claim the benefit under Title 35, United States Code §119(e) of any United States provisional application(s) listed below.

- a. ☐ no such applications have been filed.
 b. ☒ such applications have been filed as follows:

PROVISIONAL APPLICATION(S), IF ANY, UNDER 35 USC §119(c)	
APPLICATION NUMBER	DATE OF FILING (day, month, year)
60/130,338	21 April 1999

We hereby claim the benefit under Title 35, United States Code, §120 of any United States applications or §365(c) of any PCT international application(s) designating the United States of America, listed below.

- a. ☒ no such applications have been filed.
 b. ☐ such applications have been filed as follows:

APPLICATION NUMBER	DATE OF FILING (day, month, year)	STATUS (patented, pending, abandoned)

Insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code, §112, we acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application.

We hereby appoint Ann M. Muetting (Reg. No. 33,977), Kevin W. Raasch (Reg. No. 35,651), Mark J. Gebhardt (Reg. No. 35,518), Victoria A. Sandberg (Reg. No. 41,287), David L. Provence (Reg. No. 43,022), Matthew W. Adams (Reg. No. 43,459), Loren Albin (Reg. No. 37,763), Brian J. Walsh (Reg. No. 45,543), and Kathleen L. Franklin (Reg. No. 47,574) our attorneys with full powers (including the powers of appointment, substitution, and revocation) to prosecute this application and any division, continuation, continuation-in-part, reexamination or reissue thereof, and to transact all business in the Patent and Trademark Office connected therewith.

Please direct all correspondence in this case to:

Attention: Victoria A. Sandberg
 Muetting, Raasch & Gebhardt, P.A.
 P.O. Box 581415
 Minneapolis, MN 55458-1415
 Telephone No. (612) 305-
 Customer Number 26813

Declaration and Power of Attorney

Page 2 of 4

Applicant(s): Baig et al.

Serial No.: PCT/US00/10672

Int'l Filing Date: April 20, 2000

Title: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

We hereby claim the benefit under Title 35, United States Code §119(e) of any United States provisional application(s) listed below.

- a. ☐ no such applications have been filed.
b. ☒ such applications have been filed as follows:

PROVISIONAL APPLICATION(S), IF ANY, UNDER 35 USC §119(e)	
APPLICATION NUMBER	DATE OF FILING (day, month, year)
60/130,338	21 April 1999

We hereby claim the benefit under Title 35, United States Code, §120 of any United States applications or §365(c) of any PCT international application(s) designating the United States of America, listed below.

- a. ☒ no such applications have been filed.
b. ☐ such applications have been filed as follows:

APPLICATION NUMBER	DATE OF FILING (day, month, year)	STATUS (patented, pending, abandoned)

Insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code, §112, we acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application.

We hereby appoint Ann M. Mueuting (Reg. No. 33,977), Kevin W. Raasch (Reg. No. 35,651), Mark J. Gebhardt (Reg. No. 35,518), Victoria A. Sandberg (Reg. No. 41,287), David L. Provence (Reg. No. 43,022), Matthew W. Adams (Reg. No. 43,452), Loren Albin (Reg. No. 37,763), Brian J. Walsh (Reg. No. 45,543), and Kathleen L. Franklin (Reg. No. 47,574) our attorneys with full powers (including the powers of appointment, substitution, and revocation) to prosecute this application and any division, continuation, continuation-in-part, reexamination or reissue thereof, and to transact all business in the Patent and Trademark Office connected therewith.

Please direct all correspondence in this case to:

Attention: Victoria A. Sandberg
Mueuting, Raasch & Gebhardt, P.A.
P.O. Box 581413
Minneapolis, MN 55458-1415
Telephone No. (612) 305-
Customer Number 26815

Declaration and Power of Attorney

Applicant(s): Raig et al.

Serial No.: PCT/US00/10672

Int'l Filing Date: April 20, 2000

Title: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

Page 2 of 4

We hereby claim the benefit under Title 35, United States Code §119(e) of any United States provisional application(s) listed below.

- a. ☐ no such applications have been filed.
b. ☒ such applications have been filed as follows:

PROVISIONAL APPLICATION(S), IF ANY, UNDER 35 USC §119(e)	
APPLICATION NUMBER	DATE OF FILING (day, month, year)
60/130,338	21 April 1999

We hereby claim the benefit under Title 35, United States Code, §120 of any United States applications or §365(c) of any PCT International application(s) designating the United States of America, listed below.

- a. ☒ no such applications have been filed,
b. ☐ such applications have been filed as follows:

APPLICATION NUMBER	DATE OF FILING (day, month, year)	STATUS (patented, pending, abandoned)

Insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code, §112, we acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application.

We hereby appoint Ann M. Muegling (Reg. No. 33,977), Kevin W. Raasch (Reg. No. 35,651), Mark J. Gebhardt (Reg. No. 35,518), Victoria A. Sandberg (Reg. No. 41,287), David L. Provence (Reg. No. 43,022), Matthew W. Azans (Reg. No. 43,459), Loren Albin (Reg. No. 37,763), Brian J. Walsh (Reg. No. 45,543), and Kathleen L. Franklin (Reg. No. 47,574) our attorneys with full powers (including the powers of appointment, substitution, and revocation) to prosecute this application and any division, continuation, continuation-in-part, reexamination or reissue thereof, and to transact all business in the Patent and Trademark Office connected therewith.

Please direct all correspondence in this case to:

Attention: Victoria A. Sandberg
Muegling, Raasch & Gebhardt, P.A.
P.O. Box 581415
Minneapolis, MN 55458-1415
Telephone No. (612) 305-
Customer Number 26813

Declaration and Power of Attorney

Page 3 of 4

Applicant(s): Raig et al.

Serial No.: PCT/US00/10672

Int'l Filing Date: April 20, 2000

Title: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

The undersigned declare further that all statements made herein of their own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Wherefore, we pray that Letters Patent be granted to us for the invention described and claimed in the specification identified above and we hereby subscribe our names to the foregoing specification and claims, Declaration, and Power of Attorney on the date indicated below.

00 
Salma Raig

Citizenship: U.S.A.

Residence: 1907 South Millidge Avenue, Apt. E12, Athens, Georgia 30605

Mailing Address: Same

Oct 18-2001
Date

Raymond T. Damian

Citizenship: U.S.A.

Residence: 1170 Northwoods, Watkinsville, Georgia 30677

Mailing Address: Same

Date

Arthur Clinton White

Citizenship: U.S.A.

Residence: 708 Mulberry Lane, Bellaire, Texas 77401

Mailing Address: Same

Date

Declaration and Power of Attorney

Page 3 of 4

Applicant(s): Baig et al.

Serial No.: PCT/US00/10672

Int'l Filing Date: April 20, 2000

Title: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

The undersigned declare further that all statements made herein of their own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Wherefore, we pray that Letters Patent be granted to us for the invention described and claimed in the specification identified above and we hereby subscribe our names to the foregoing specification and claims, Declaration, and Power of Attorney on the date indicated below.

Salman Baig

Date

Citizenship: U.S.A.

Residence: 1907 South Milledge Avenue, Apt. B12, Athens, Georgia 30605

Mailing Address: Same

280 Raymond V. Dumian

10/17/01

Date

Citizenship: U.S.A.

Residence: 1170 Northwoods, Watkinsville, Georgia 30677

Mailing Address: Same GA

Arthur Clinton White

Date

Citizenship: U.S.A.

Residence: 708 Mulberry Lane, Bellaire, Texas 77401

Mailing Address: Same

Declaration and Power of Attorney

Page 3 of 4

Applicant(s): Baig et al.

Serial No.: PCT/US00/10672

Int'l Filing Date: April 20, 2000

Title: CYSTEINE PROTEASE AND INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

The undersigned declare further that all statements made herein of their own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Wherefore, we pray that Letters Patent be granted to us for the invention described and claimed in the specification identified above and we hereby subscribe our names to the foregoing specification and claims, Declaration, and Power of Attorney on the date indicated below.

Salman Baig

Date

Citizenship: U.S.A.

Residence: 1907 South Milledge Avenue, Apt. E12, Athens, Georgia 30605

Mailing Address: Same

Raymond T. Damian

Date

Citizenship: U.S.A.

Residence: 1170 Northwoods, Watkinsville, Georgia 30677

Mailing Address: Same

300 Arthur Clinton White

Date

Citizenship: U.S.A.

Residence: 708 Mulberry Lane, Bellaire, Texas 77401

Mailing Address: Same

TX

10/17/01

§ 1.56 Duty to disclose information material to patentability.

(a) A patent by its very nature is affected with a public interest. The public interest is best served, and the most effective patent examination occurs when, at the time an application is being examined, the Office is aware of and evaluates the teachings of all information material to patentability. Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is cancelled or withdrawn from consideration, or the application becomes abandoned. Information material to the patentability of a claim that is cancelled or withdrawn from consideration need not be submitted if the information is not material to the patentability of any claim remaining under consideration in the application. There is no duty to submit information which is not material to the patentability of any existing claim. The duty to disclose all information known to be material to patentability is deemed to be satisfied if all information known to be material to patentability of any claim issued in a patent was cited by the Office or submitted to the Office in the manner prescribed by §§ 1.97(b)-(d) and 1.98. However, no patent will be granted on an application in connection with which fraud on the Office was practiced or attempted or the duty of disclosure was violated through bad faith or intentional misconduct. The Office encourages applicants to carefully examine:

- (1) Prior art cited in search reports of a foreign patent office in a counterpart application, and
- (2) The closest information over which individuals associated with the filing or prosecution of a patent application believe any pending claim patentably defines, to make sure that any material information contained therein is disclosed to the Office.

(b) Under this section, information is material to patentability when it is not cumulative to information already of record or being made of record in the application, and

- (1) It establishes, by itself or in combination with other information, a prima facie case of unpatentability of a claim; or
- (2) It refutes, or is inconsistent with, a position the applicant takes in:
 - (i) Opposing an argument of unpatentability relied on by the Office, or
 - (ii) Asserting an argument of patentability.

A prima facie case of unpatentability is established when the information compels a conclusion that a claim is unpatentable under the preponderance of evidence, burden-of-proof standard, giving each term in the claim its broadest reasonable construction consistent with the specification, and before any consideration is given to evidence which may be submitted in an attempt to establish a contrary conclusion of patentability.

(c) Individuals associated with the filing or prosecution of a patent application within the meaning of this section are:

- (1) Each inventor named in the application;
- (2) Each attorney or agent who prepares or prosecutes the application; and
- (3) Every other person who is substantively involved in the preparation or prosecution of the application and who is associated with the inventor, with the assignee or with anyone to whom there is an obligation to assign the application.

(d) Individuals other than the attorney, agent or inventor may comply with this section by disclosing information to the attorney, agent, or inventor.

(e) In any continuation-in-part application, the duty under this section includes the duty to disclose to the

Office all information known to the person to be material to patentability, as defined in paragraph (b) of this section, which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

Applicant(s): Baig et al.

Serial No.: PCT/US00/10672

Int'l Filing Date: April 20, 2000

Title: CYSTEINE PROTEASE INHIBITORS FOR PREVENTION AND TREATMENT OF NEUROCYSTICERCOSIS

§ 1.56 Duty to disclose information material to patentability.

(a) A patent by its very nature is affected with a public interest. The public interest is best served, and the most effective patent examination occurs when, at the time an application is being examined, the Office is aware of and evaluates the teachings of all information material to patentability. Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is cancelled or withdrawn from consideration, or the application becomes abandoned. Information material to the patentability of a claim that is cancelled or withdrawn from consideration need not be submitted if the information is not material to the patentability of any claim remaining under consideration in the application. There is no duty to submit information which is not material to the patentability of any existing claim. The duty to disclose all information known to be material to patentability is deemed to be satisfied if all information known to be material to patentability of any claim issued in a patent was cited by the Office or submitted to the Office in the manner prescribed by §§ 1.97(b)-(d) and 1.98. However, no patent will be granted on an application in connection with which fraud on the Office was practiced or attempted or the duty of disclosure was violated through bad faith or intentional misconduct. The Office encourages applicants to carefully examine:

- (1) Prior art cited in search reports of a foreign patent office in a counterpart application; and
- (2) The closest information over which individuals associated with the filing or prosecution of a patent application believe any pending claim patentably defines, to make sure that any material information contained therein is disclosed to the Office.

(b) Under this section, information is material to patentability when it is not cumulative to information already of record or being made of record in the application, and

- (1) It establishes, by itself or in combination with other information, a *prima facie* case of unpatentability of a claim; or
- (2) It refutes, or is inconsistent with, a position the applicant takes in:
 - (i) Opposing an argument of unpatentability relied on by the Office; or
 - (ii) Asserting an argument of patentability.

A *prima facie* case of unpatentability is established when the information compels a conclusion that a claim is unpatentable under the preponderance of evidence, burden-of-proof standard, giving each term in the claim its broadest reasonable construction consistent with the specification, and before any consideration is given to evidence which may be submitted in an attempt to establish a contrary conclusion of patentability.

(c) Individuals associated with the filing or prosecution of a patent application within the meaning of this section are:

- (1) Each inventor named in the application;
- (2) Each attorney or agent who prepares or prosecutes the application; and
- (3) Every other person who is substantively involved in the preparation or prosecution of the application and who is associated with the inventor, with the assignee or with anyone to whom there is an obligation to assign the application.

(d) Individuals other than the attorney, agent or inventor may comply with this section by disclosing information to the attorney, agent, or inventor.

(e) In any continuation-in-part application, the duty under this section includes the duty to disclose to the

Office all information known to the person to be material to patentability, as defined in paragraph (b) of this section, which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.